



Influence of Key Health Related Indicators on Adult Mortality: Result from UN Member Countries

*Chhabi Lal RANABHAT^{1,2,3}, Myung-Bae PARK⁴, *Chun-Bae KIM^{1,5}, Chang-Soo KIM^{1,6}, Hyoung-Sun JEONG⁷, Sang Baek KOH⁵, Sei-Jin CHANG⁵

1. Institute for Poverty Alleviation and International Development, Yonsei University, Wonju, Gangwon-do, Korea
2. Health Science Foundations and Study Center, GPO-44600, Kathmandu, Nepal
3. Manmohan Institute of Health Science, Kathmandu, Nepal
4. Dept. of Gerontal Health and Welfare, Pai Chai University, College of Howard, Daejeon, Korea
5. Dept. of Preventive Medicine, Wonju College of Medicine, Yonsei University, Wonju, Gangwon-do, Korea
6. Dept. of Business Administration, College of Government and Business, Yonsei University, Wonju, Gangwon-do, Korea
7. Dept. of Health Administration, Yonsei University, Wonju, Gangwon-do, Korea

*Corresponding Authors: Emails: chhabir@gmail.com, kimcb@yonsei.ac.kr

(Received 15 Apr 2017; accepted 19 Sep 2017)

Abstract

Background: Adult mortality is associated with different demographic and behavioral risk factors including approaches to health care financing. Adult mortality rate significantly reflects the effectiveness of public health-related program and intervention. The aim of this study was to find strength of association between key health's related indicators and adult mortality rate.

Methods: This cross-sectional study used 5 sets of data combined into one from different organizations of 193 countries using record linkage theory. Eleven key health-related indicators were taken as independent variables and adult mortality of male and female were dependent variables from 2010 to 2013. Average mortality for male and female was shown by means and standard deviations, raw association by Pearson correlation and strength of association by hierarchical linear regression.

Results: The average adult mortality rate (AMR) of male was 0.209 ± 0.106 and of female, 0.146 ± 0.105 in years. In raw correlation, almost all health indicators were associated with AMR of male and female. In regression analysis, Universal Health Coverage (UHC) significantly reduced (male ~ 0.43 , female ~ 0.30) adult mortality, in contrast, population growth significantly increased (male ~ 0.37 , female ~ 0.43). Alcohol consumption per year increased AMR in male by 0.41 ($P < 0.01$) and vaccination coverage (DPT 3) significantly reduced the AMR (0.26) in female.

Conclusion: It is necessary to extend the UHC in remaining countries and still a need to control the population where there is high population growth. Effectively control of alcoholic drink in male and full coverage of vaccination in childhood mitigates adult mortality. The UHC is ambitious goal for SDG and special attention should be provided nationally and globally

Keywords: Adult mortality rate, Health indicators, Universal health coverage, Record linkage

Introduction

Adult deaths threaten the livelihood of entire families (1) and seriously affect the children's de-

velopment (2,3) limit support on elderly (4,5) and economic activities of households (6). There is

massive inequality in adult mortality. In some countries, the adult mortality is even higher after 2010 than that of 1990 and by sex, there is the highest women mortality in South Asia (7). Lesotho has the highest (0.50) and San Marino has the lowest (0.05) male adult mortality rate which is 10 times gap each other (8). Similarly, female adult mortality rate is 23 times higher in Lesotho (0.50) in comparison with Honduras (0.02) (9). South African experience shows that adults could be protected with healthy lifestyle and behavior, disease prevention approaches and adult friendly health policy (10). There are many factors associated adult mortality, which is different from neonatal, infant and maternal mortality. The risk factors; injuries, lifestyle, sexual behavior and other demographic factors were explored (11) but the situation of adult mortality with different varieties of health indicators have seldom been discussed. The concept of adult mortality rate was developed in Europe to investigate the economic value of human in productive age (12). In comparison with all mortality, adult mortality is more important in term of national economy, sustainable development and family protection (13) and adult mortality is higher than other mortality due to more exposure with life-threatening events and injuries (14). Studies in adult mortality rate; (important health indicator) are not enough and the relation with related programs and indicators are extremely dearth. Health status, risk and outcome indicators are necessary to set the priority and long-term health planning (15) and WHO highlighted that health indicators have significant role in sustainable planning and development (16). A family health survey in India suggested that adult behavioral indicators (17) and sanitation indicators with infectious disease (18) are associated with adult mortality. To reduce adult death, it is necessary formulate special policy but before this, it is obligatory to find out the association between health-related indicator and adult mortality. Global, regional and national adult mortality show that infectious diseases, major noncommunicable diseases like heart problems, cancer, injuries, poor immunity in childhood and adapted health care system have strong influences

on adult mortality (19-21). Hygiene and sanitation, alcohol and tobacco use, child vaccination and health care expenditure explicitly determine the adult mortality. Socio-behavioral factors and biological factors (22,23) were risk factors for adult death. Education, which is the means for life, affects the mortality and morbidity of adults (24,25). The structural factors like social conflicts, economic disparities, and health financing police are strongly associated with adult mortality (26). Health expenditures are strongly associated with health outcomes and especially with reduction in child and adult mortality (27). The association between health indicators and quality of life was explored (28) however there was not a lot of research using that model.

The study and analysis of adult mortality tend to focus on risk factors and disease burden. In other words, research on adult mortality has been limited to individual factors but indifferent to program and indicators. There could be multiple associations between different health indicators like social determinants, human development and universal health coverage (UHC) (29) so that it is possible to access the proxy effect among them. This study will observe the association between demographics, diseases prevention, lifestyle and health financing indicators with adult mortality in a global context among UN country members. Probably, it is a first study to observe the vital health-related indicators for adult mortality and using data from diverse sources in different periods.

Therefore, the aim of this study was to find the strength of association between key health-related indicator and adult mortality from UN country members.

Materials and Methods

Study design

In this multi-country cross-sectional study variables from different organizations and timeline were used. The outcome variable, adult mortality rate was for the 2012-13 period and input variables prior to 2013.

Data adoption model and sources

Data were created based on the Record Linkage theory applied by Dunn HL in 1946 (30). The objectives of this model were to preserve the records, verify them and create the new statistics, however, those data originated in different purpose. Furthermore, this model was advanced by Jutte DP 2011 as Administrative Record Linkage as a tool for public health research (31). We made the database from World Bank health data 2013 (32), WHO health report tobacco epidemic (33) Full complete data set

and alcohol information (34) and “The political economy of UHC (35) (Table 1).

Data management

We created new data by country for all 193 UN members. Data and outliers were cross-checked and missing data from some countries were excluded from the study. After verification and re-verification, the data were exported into SPSS -20 version (Chicago, IL, USA).

Table 1: Source of database

S.N	Data sources	Reference	Full complete data set
1	World Bank health data	World Bank: Data Catalogue	
2	Category of countries adopted and achieved universal health coverage	Stuckler D et.al The political economy of universal health coverage; 2010	
3	WHO health report tobacco epidemic 2013	World Health Report 2014	
4	WHO alcohol information report 2010	World Health Report 2014	

Data analysis model

1. The descriptive results were presented with means and standard deviations as global average.
2. In the second phase, Pearson correlation coefficient was computed.
3. In third step, multi-variant analysis was performed using hierarchal linear regression models

The data were interpreted by beta coefficient and *P*-value at the level of <0.05 and <0.001. The conclusion was drawn from the results hierarchical linear regression full model.

Structure of variables

All of the above variables are numerical except universal health coverage. UHC is combined with 3 indicators; legal obligation of UHC, skill birth attendance rate and health insurance coverage. Therefore, we applied the UHC as yes =1 and no =0 category. After making consistent variables, we performed the linear regression models.

Validity and reliability

We verified Word Bank, WHO and UNDP data used for this study. Likewise, the normality of

data was checked by the observation of histogram and consistency of data was checked by Cronbach's alpha for appropriate variables. Universal health coverage is very wide concept and needs to check multicollinearity. Multicollinearity was checked by variance inflation factors (VIF) with the standard of less than 3.

Ethical aspects

The data is open access to everyone from above organizations and no need to take ethical approval.

Results

Descriptive statistics

Fifty-eight countries (30%) have achieved universal health care. Primary school enrollment and child vaccination (DPT- 3) was almost achieved (>85%). Average population growth was less than 2%, more than 1/3rd population did not achieve sanitation coverage and out-of-pocket payment for treatment was 61.1%. The average adult mortality rate of both sexes was 0.167 ± 0.108 and male and female AMR was 0.209 ± 0.106 and 0.146 ± 0.105 (Table 2).

Table 2: Descriptive status of all health indicators

<i>Variables (in years)</i>	<i>No. of countries</i>	<i>Mean ± SD</i>
Net primary school enrollment (NPSE - 012) %	147	95.8±31.0
Economic Growth Rate (EGR 06 - 010)%	177	3.4±2.3
Population Growth Rate (PGR - 012)%	191	1.4±1.3
Child Vaccination Coverage (DPT 3 - 012)%	189	89.2±13.0
Sanitation Coverage (SAN COV – 012)%	193	61.2±27.3
Average adult consumption (ALCOHOL CON - 010) in liter/year	174	5.9±3.9
Prevalence of adult smoking - male (Prev. AS male - 012) %	134	32.3±12.6
Prevalence of adult smoking -female (Prev. AS female - 012) %	134	11.2±10.4
Out of Pocket Payment (OPP - 012) %	185	61.1±39.5
Total Health Expenditure (THE - 012) %	184	6.7±2.9
Government Health Expenditure per year (GHE - 012) %	186	11.6±4.9
Adult mortality both sex (012)/thousand/year	181	167.1±108.8
Adult mortality rate-male (012) / thousand/year	181	209.9±106.1
Adult mortality rate-female (012) / thousand/year	181	146.3±105.1
Achievement of Universal Health Care (UHC) –Yes (N ₁ -193)	58	30%

Cronbach's alpha 0.712, Variance inflation factors (VIF) with UHC and other factors <1.5

Table 3 shows the correlation status of all variables. Adult male mortality is positively associated with population growth rate, alcohol consumption per year, prevalence adult smoking, out of pocket payment and government health expenditure ($P<0.01$). Likewise, it is negatively associated with economic growth rate, DPT 3 and sanitation coverage and universal health coverage. Similar trend can be observed in female adult mortality

status. Female adult mortality had a strong positive association with population growth and strong negative association with DPT 3, sanitation and universal health coverage.

Adult mortality by hierarchical regression model

Table 4 shows the association between all predicted indicators and male adult mortality.

Table 3: Situation of raw correlation among all variables using Pearson’s correlation (*P-value<0.05, **P<0.01)

<i>S. N</i>	<i>Variables</i>	<i>1</i>	<i>2</i>	<i>3</i>	<i>4</i>	<i>5</i>	<i>6</i>	<i>7</i>	<i>8</i>	<i>9</i>	<i>10</i>	<i>11</i>	<i>12</i>	<i>13</i>	<i>14</i>
1	NPSE (012)	1													
2	EGR (06 - 010)	.213*	1												
3	PGR (012)	-.109	.349**	1											
4	DPT3 (012)	.046	-.146	-.335**	1										
5	SAN COV (012)	.065	.021	.150	.294**	1									
6	ALCOHOL CON	-.240**	-.282**	.194	.192	-.029	1								
7	Prev. AS – male (012)	-.048	.023	.523**	.131	.152*	.533**	1							
8	Prev. AS - female (012)	-.078	-.411**	.399**	.102	.151	0.24	.318**	1						
9	UHC	.003	.241**	-.272**	.292**	.147*	-.137	-.107	-.505**	1					
10	OPP (012)	-.074	-.204**	.358**	.114	.053	.381**	.037	.502**	-.202**	1				
11	THE (012)	.028	-.051	-.051	.030	.011	.071	.023	.082	.098	.116	1			
12	GHE (012)	.003	-.128	-.273**	.205**	.101	-.245**	-.079	-.359*	.240**	.598**	-.101	1		
13	AMR- male (012)	-.031	-.250*	.533**	-.480**	-.492**	.575**	.407**	.180	-.597**	.351**	.085	.336**	1	
14	AMR- female (012)	-.010	-.231**	.498**	-.514**	-.467**	.168	.258*	.304**	-.554*	.294*	.055	.271*	.608**	1

For the full model (model 4), AMR would significantly increase by 0.379 and 0.419 with population growth rate ($P < 0.05$) and alcohol consumption per year. In contrast, AMR would significantly decrease ($P < 0.01$, $\beta \sim 0.433$) with universal health coverage.

In regression, the determinants health indicators of female adult mortality were little different.

Lifestyle indicators were not associated with the full model but in model 3, female smoking prevalence was positively associated with mortality ($P < 0.05$). In model 4 (full), female adult mortality was significantly increased ($P < 0.01$, $\beta \sim 0.437$) with population growth rate and significantly decreased with DPT 3 coverage ($\beta \sim 0.262$) and universal health coverage ($\beta \sim 0.301$) (Table 5).

Table 4: Linear regression among demographic, disease prevention, lifestyle, health financing indicators and male adult mortality rate

Predictors	Adult mortality rate male (012)							
	Model 1		Model 2		Model 3		Model 4 (full)	
	St beta Coeff.	Sig.	St Beta Coeff.	Sig.	St Beta Coeff.	Sig.	St Beta Coeff.	Sig.
Eco-demographic indicators								
NPSE (012)	-0.043	0.597	-0.010	0.894	0.068	0.479	0.100	0.265
EGR (06- 010)	0.082	0.396	0.111	0.236	0.141	0.209	0.064	0.561
PGR (012)	0.499	<0.001	0.380	<0.001	0.475	0.004	0.379	0.013
Disease prevention indicators								
DPT3 (012)			-0.265	0.003	-0.199	0.069	-0.154	0.120
SAN COV (012)			-0.131	0.115	-0.188	0.066	-0.096	0.298
Lifestyle indicators								
Adult smoking male					0.099	0.368	0.079	0.429
ALCOHOL CON					0.279	0.034	0.419	0.002
Health financing indicators								
UHC (Yes/No)							-0.433	<0.001
OPP (012)							-0.108	0.358
THE (012)							-0.039	0.648
GHE (012)							-0.037	0.723
R ²	0.301		0.323		0.346		0.517	
Adj R ²	0.274		0.275		0.283		0.448	
F value	10.355		9.846		5.776		6.912	
P-value	<0.001		<0.001		<0.001		<0.001	

Note: The details of variables have been presented in methodology and Table 2.

Table 5: Linear regression among demographic, disease prevention, lifestyle, health financing indicators and female adult mortality rate

Predictors	Adult mortality rate female (012)							
	Model 1		Model 2		Model 3		Model 4 (full)	
	St Beta Coeff.	Sig.	St Beta Coeff.	Sig.	St Beta Coeff.	Sig.	St Beta Coeff.	Sig.
Eco-demographic indicators								
NPSE (012)	-0.076	0.314	-0.046	0.521	0.094	0.260	0.078	0.352
EGR (06 - 010)	0.018	0.839	0.035	0.680	-0.064	0.503	-0.086	0.398
PGR (012)	0.586	0.000	0.463	0.000	0.490	0.000	0.437	0.001
Disease prevention indicators								
DPT3 (012)			-0.223	0.006	-0.273	0.005	-0.262	0.008
SAN COV (012)			-0.188	0.014	-0.134	0.129	-0.106	0.224
Lifestyle indicators								
Adult smoking female					0.242	0.037	0.171	0.166
ALCOHOL CON					0.198	0.069	0.188	0.126
Health financing indicators								
UHC (Yes/No)							-0.301	0.003
OPP (012)							0.127	0.270
THE (012)							-0.024	0.758
GHE (012)							-0.045	0.634
R ²	0.331		0.429		0.521		0.576	
Adj R ²	0.334		0.404		0.477		0.511	
F value	19.721		17.525		11.965		8.848	
P value	<0.001		<0.001		<0.001		<0.001	

Note: The details of variables have been presented in methodology and Table 2.

Discussion

Adult mortality is directly related to disease or condition of the individual but adult mortality rate is a key health indicator and associated with other related indexes. The average male adult mortality was 1.43 times higher than females (Table 2). In the US (36) and Japan (37), due to several risk factors for diseases and injury, adult mortality was higher in men. In raw correlation, adult mortality of both sexes was correlated with almost independent variables however, the result was interpreted by linear regression full model (model 4). Population growth rate, alcohol consumption, universal health coverage and vaccine coverage (DPT 3) childhood are strongly associated ($P<0.05$) with adult mortality rate among 11 major indicators in linear regression (full model). In other words, this study is more useful to observe related program impact on adult mortality.

There is a linear positive association between population growth rate and adult mortality for men and women. The adult mortality of men and women would increase by 0.379 and 0.437 with population growth rate. This estimation is similar to the demographic transition formula (38) and population estimation (39). The adult mortality is high in South Asia, Africa and Middle East Asia where population growth is also high (40). Similarly, there is linear negative association between universal health care and adult mortality status (26). The adult mortality would decrease by 0.433 and 0.301 respectively for men and women with universal health coverage as like above study. UHC overall improves the population health by equity and access of preventive, promotive, curative and palliative health care without financial constraints (41) and health program like vaccination, sanitation, maternal and child health, nutrition, health education etc. are assured so that the adults could be strong from childhood. As a health care service modality, UHC with universal cooperative health insurance would reduce adult death (42).

Among men, the adult mortality is significantly higher ($P=0.002$, $\beta\sim 0.41$) with alcohol consumption. There is no established causal relationship of alcohol consumption with fatal diseases but

male drinking behavior would be responsible. Usually, male adult mortality increases with heavy and uncontrolled drinking behavior (43,44) and this behavior has multiple impacts; increasing violence and murder, worsening of infectious diseases like tuberculosis and hepatitis, limiting life expectancy with liver and heart diseases, increasing mental diseases and suicide and increasing unintentional injuries due to drunk driving (45). Moreover, unprotected sex behavior is common after alcohol consumption and the risk for sexually transmitted diseases particularly HIV is increased (46). In South Africa, alcohol users were more vulnerable to risky sexual intercourse (47) and the prevalence of adult mortality was high in Africa due to this circumstance.

The determinant indicators for male and female adult mortality are different. Child vaccination DPT -3 would reduce female adult mortality by 0.262 ($P<0.01$). The girl's death rate was 2 times higher for those not received the DPT vaccine in comparison with those who received the vaccine (48) which is similar to our findings. However, DPT vaccine was associated with higher mortality for girls (49) and obviously could not protect through the adult stage. The coverage of this study was wide and included all UN country members and 5 different sources of data. Major determinant indicators of adult mortality rate were examined however, there were not equal samples for all variables examined and those countries numbers may not enough from statistical perspective.

Conclusion

Population growth rate and alcohol consumption amount would increase the adult mortality rate but it can be reduced by universal health coverage. The UHC is ambitious goal for SDG and special attention should be provided nationally and globally.

Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or fal-

sification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Acknowledgements

This research is supported by Korean Medical Association (RIHP-2015-02) and National Research Foundation of Korea (NRF-2016S1A5B892520). We also thank Editage for editing the English content of this paper.

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. Yamano T, Jayne TS (2004). Measuring the impacts of working-age adult mortality on small-scale farm households in Kenya. *World Development*, 32(1): 91-119.
2. Case A, Ardington C (2006). The impact of parental death on school outcomes: Longitudinal evidence from South Africa. *Demography*, 43(3): 401-20.
3. Evans DK, Miguel E (2007). Orphans and schooling in Africa: A longitudinal analysis. *Demography*, 44(1): 35-57.
4. Ainsworth M, Dayton J (2003). The impact of the AIDS epidemic on the health of older persons in northwestern Tanzania. *World Development*, 31(1): 131-48.
5. Kautz T, Bendavid E, Bhattacharya J, Miller G (2010). AIDS and declining support for dependent elderly people in Africa: Retrospective analysis using demographic and health surveys. *BMJ*, 340: c2841:1-6.
6. Chao L-W, Pauly M, Szrek H et al (2007). Poor health kills small business: Illness and microenterprises in South Africa. *Health Aff (Millwood)*, 26(2): 474-82.
7. Rajaratnam JK, Marcus JR, Levin-Rector A et al (2010). Worldwide mortality in men and women aged 15–59 years from 1970 to 2010: A systematic analysis. *Lancet*, 375(9727): 1704-20.
8. World Bank (2013). Mortality rate, adult, male (per 1,000 male adults). <http://data.worldbank.org/indicator/SP.DY.N.AMRT.MA>
9. World Bank (2013). Mortality rate, adult, female (per 1,000 female adults). <http://data.worldbank.org/indicator/SP.DY.N.AMRT.FE>
10. Sartorius B, Kahn K, Collinson MA et al (2013). Dying in their prime: Determinants and space-time risk of adult mortality in rural South Africa. *Geospat Health*, 7(2): 237-49.
11. Gregg EW, Cheng YJ, Saydah S et al (2012). Trends in death rates among US adults with and without diabetes between 1997 and 2006: Findings from the national health interview survey. *Diabetes Care*, 35(6): 1252-7.
12. Hill K (2003). Adult mortality in the developing world: What we know and how we know it: UN. <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.414.7365&rep=rep1&type=pdf>
13. Rogers RG, Crimmins EM (2011). *International Handbook of Adult Mortality*: Springer Science & Business Media.
14. Norman R, Matzopoulos R, Groenewald P, Bradshaw D (2007). The high burden of injuries in South Africa. *Bull World Health Organ*, 85(9):695-702.
15. Pilav A (1999). Health status indicators and the importance of their use in daily practice. *Med Arh*, 53(3 Suppl 3):51-3.
16. Schirmding Yv (2002). *Health in Sustainable Development Planning: The Role of Indicators*. Geneva: World Health Organization.
17. Saikia N, Bhat PM (2008). Factors affecting adult mortality in India: An analysis of National Family Health Surveys of 1992-93 and 1998-99 (NFHS I and II). *Demography India*, 37(2): 291-302.
18. Ram U, Jha P, Gerland P et al (2015). Age-specific and sex-specific adult mortality risk in India in 2014: Analysis of 0.27 million nationally surveyed deaths and demographic estimates from 597 districts. *Lancet Glob Health*, 3(12): e767-75.
19. Naghavi M, Wang H, Lozano R et al (2015). Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: A systematic

- analysis for the Global Burden of Disease Study 2013. *Lancet*, 385(9963): 117-71.
20. Misganaw A, Mariam DH, Araya T (2013). Association of socioeconomic and behavioral factors with adult mortality: Analysis of data from verbal autopsy in Addis Ababa, Ethiopia. *BMC Public Health*, 13: 634-45.
 21. Setel P, Kitange H, Alberti K, Moshiro C (1998). The policy implications of adult morbidity and mortality in Tanzania: From data analysis to health policy—preliminary experiences. Global Forum for Health Research (Forum 2). https://research.ncl.ac.uk/ammp/site_files/public_html/geneva.pdf
 22. Pensola TH, Martikainen P (2003). Cumulative social class and mortality from various causes of adult men. *J Epidemiol Community Health*, 57(9): 745-51.
 23. Rogers RG, Everett BG, Saint Onge JM, Krueger PM (2010). Social, behavioral, and biological factors, and sex differences in mortality. *Demography*, 47(3): 555-78.
 24. Bostock S, Steptoe A (2012). Association between low functional health literacy and mortality in older adults: Longitudinal cohort study. *BMJ*, 344: e1602:1-10.
 25. Clark D, Roayer H (2013). The effect of education on adult mortality and health: Evidence from Britain. *Am Econ Rev*, 103(6): 2087-120.
 26. Ranabhat CL, Kim CB, Park MB, Acharaya S (2017). Multiple disparities in adult mortality in relation to social and health care perspective: Results from different data sources. *Global Health*, 13(1): 57-65.
 27. Moreno-Serra R, Smith P (2011). The effects of health coverage on population outcomes: a country-level panel data analysis. Washington, DC: Results for Development Institute.
 28. Schmidt S, Power M, Bullinger M, Nosikov A (2005). The conceptual relationship between health indicators and quality of life: results from the cross-cultural analysis of the EUROHIS field study. *Clin Psychol Psychother*, 12(1): 28-49.
 29. De Andrade LOM, Filho AP, Solar O et al (2015). Social determinants of health, universal health coverage, and sustainable development: Case studies from Latin American countries. *Lancet*, 385(9975): 1343-51.
 30. Dunn HL (1946). Record Linkage. *Am J Public Health Nations Health*, 36(12): 1412-6.
 31. Jutte DP, Roos LL, Brownell MD (2011). Administrative record linkage as a tool for public health research. *Annu Rev Public Health*, 32: 91-108.
 32. World Bank (2015). Data. <https://data.worldbank.org/>
 33. World Health Organization (2014). *Global Health Observatory (GHO)*. World Health Organization.
 34. World Health Organization (2010). *WHO Global Information System on Alcohol and Health (GISAH)*. Geneva. <http://www.who.int/gho/alcohol/en/>
 35. Stuckler D, Feigl AB, Basu S, McKee M (2010). *The Political Economy of Universal Health Coverage*. Background paper for the global symposium on health systems research, World Health Organization, Geneva.
 36. Yang YC, McClintock MK, Kozloski M, Li T (2013). Social isolation and adult mortality: The role of chronic inflammation and sex differences. *J Health Soc Behav*, 54(2): 183-203.
 37. Suzuki E, Kashima S, Kawachi I, Subramanian S (2012). Social and geographic inequalities in premature adult mortality in Japan: A multilevel observational study from 1970 to 2005. *BMJ Open*, 2(2): e000425:1-11.
 38. Galor O (2012). The demographic transition: Causes and consequences. *Cliometrica (Berl)*, 6(1): 1-28.
 39. Bongaarts J (2009). Human population growth and the demographic transition. *Philos Trans R Soc Lond B Biol Sci*, 364(1532): 2985-90.
 40. Gribble JN (2012). *World Population Data Sheet 2012*. Washington, DC: Population Reference Bureau.
 41. Moreno-Serra R, Smith PC (2012). Does progress towards universal health coverage improve population health? *Lancet*, 380(9845): 917-23.
 42. Ranabhat CL, Kim CB, Singh DR, Park MB (2017). A comparative study on outcome of government and co-operative community-based health insurance in Nepal. *Front Public Health*, 5: 250:1-9.
 43. Plunk AD, Syed- Mohammed H, Cavazos-Rehg P et al (2014). Alcohol consumption,

- heavy drinking, and mortality: Rethinking the J- Shaped curve. *Alcohol Clin Exp Res*, 38(2): 471-8.
44. Ranabhat CL, Kim CB, Park MB, Bajgai (2018) J. Impact of Spiritual Behavior on Self-Reported Illness: A Cross-Sectional Study among Women in the Kailali District of Nepal. *J of Lifestyle Med*. 8(1):23.
45. Rehm J, Baliunas D, Borges GL et al (2010). The relation between different dimensions of alcohol consumption and burden of disease: An overview. *Addiction*, 105(5): 817-43.
46. Majer JM, Rodriguez J, Bloomer C, Jason LA (2014). Predictors of HIV-risk sexual behavior examining lifetime sexual and physical abuse histories in relation to substance use and psychiatric problem severity among ex-offenders. *J Am Psychiatr Nurses Assoc*, 20(2): 138-46.
47. Burnhams NH, Parry C, Laubscher R, London L (2014). Prevalence and predictors of problematic alcohol use, risky sexual practices and other negative consequences associated with alcohol use among safety and security employees in the Western Cape, South Africa. *Subst Abuse Treat Prev Policy*, 9: 14:1-10.
48. Aaby P, Ravn H, Roth A et al (2012). Early diphtheria-tetanus-pertussis vaccination associated with higher female mortality and no difference in male mortality in a cohort of low birthweight children: An observational study within a randomised trial. *Arch Dis Child*, 97(8): 685-91.
49. Aaby P, Benn C, Nielsen J et al (2012). Testing the hypothesis that diphtheria-tetanus-pertussis vaccine has negative non-specific and sex-differential effects on child survival in high-mortality countries. *BMJ Open*, 2(3): e000707.