



# Human Health Risk Assessment of Cd, Cu, Pb and Zn through Consumption of Raw and Pasteurized Cow's Milk

*\*Soheil SOBHANARDAKANI*

*Dept. of the Environment, College of Basic Sciences, Hamedan Branch, Islamic Azad University, Hamedan, Iran*

**\*Correspondence:** Email: s\_sobhan@iauh.ac.ir

(Received 16 Feb 2017; accepted 09 Jun 2017)

## **Abstract**

**Background:** The analysis of the residual contents of heavy and toxic metals in foodstuff especially milk could be an important indicator of safety, quality, and level of pollution of the region in which the milk was produced. Therefore, the present investigation was carried out to assess human health risk of residues levels of four metals (Cd, Cu, Pb, and Zn) through consumption of raw and pasteurized cow's milk.

**Methods:** In this analytical-observational study, totally 72 samples of raw and pasteurized cow's milk samples were collected from market basket of Hamadan City, western Iran in 2014. After preparation and processing the samples in the laboratory, the concentration of metals were determined using inductively coupled plasma (ICP-OES). Moreover, all statistical analyses were performed using the SPSS statistical package according to Shapiro-Wilk test for normality, One Way ANOVA (Duncan Multiple Range Test), Independent t-test and Pearson's correlations.

**Results:** The mean concentrations ( $\mu\text{g}/\text{kg}$ ) of Cd, Cu, Pb and Zn in raw milk samples were  $0.36 \pm 0.28$ ,  $9.77 \pm 3.91$ ,  $32.83 \pm 20.80$  and  $253.70 \pm 87.96$ , respectively and in the pasteurized milk samples were  $5.57 \pm 9.33$ ,  $8.41 \pm 5.99$ ,  $25.54 \pm 26.50$  and  $90.12 \pm 91.52$ , respectively. HRI values in adults and children via consumption of raw and pasteurized cow's milk were within the safe limits ( $\text{HRI} < 1$ ).

**Conclusion:** Considering the serious contamination of some samples of raw and pasteurized milk by Cd, Pb and Zn, a control of heavy metals content during the whole production processing of milk must be applied.

**Keywords:** Food safety, Metal contamination, Health risk index, Milk

## **Introduction**

The rapid development of our society in the past few decades, especially with priority of industrial activities, mining, fuel combustion, development of mechanized cultivation and consequently the use of large amount of agricultural inputs without attention to the their environmental and health impact, caused a wide distribution of heavy metals and other hazardous pollutants in the environment and finally in the trophic chain (1, 2).

The products with animal origin specifically milk and dairy products play an important role in hu-

man diet and acquire a special significance in infant nutrition considering their protein and mineral contents essential (1). Therefore, entry of high levels of heavy metals into the dairy products could be a potential risk, serious diseases and public health problems (3-5) and for this reason the determination of the heavy metals contents in livestock's milk could be an important indicator of the hygienic condition of this product and also the level of pollution of the area in which the milk was produced (3, 6, 7).

Cadmium is toxic and non-essential for human health and accumulates principally in the kidneys and liver. Metal industries and sewers are known as the most important sources of Cd pollution (8).

Copper is one of most abundant trace elements with vitamin-like impact in human body and living systems and found in a wide range of foods we eat such as nuts, fruits, vegetables, red meat and shellfish (9, 10).

Lead can enter the food chain through contaminated soil. Lead poisoning can affect the blood circulation, kidneys, and peripheral and central nervous systems especially in adults (11). Furthermore, infants and children are at a particularly high risk for neurotoxic and developmental disorders due to this toxic element (12, 13).

However, zinc plays an important role in cancer etiology and harms some physiological activities, but this element known as a functional and structural element for human health (14, 15).

The human health risk assessment requires identification, collection, and integration of information on the chemicals health hazards, exposure of human to the chemical and relationships between exposure, dose and adverse effects (16). In this regard, a human potential health risk assessment is the process to estimate the nature and possibility of adverse health effects in humans exposed to toxins and chemicals in polluted environmental media, now or in the future (17).

The survey of potential health risks is done by compared the acceptable daily intake (ADI) of trace elements with the provisional tolerable weekly intake (PTWI) or oral reference dose (RfD). According to the standard methods (USEPA), the risk of chronic-toxic effects (health risk index (HRI) or target hazard quotients (THQ)) is described as the ratio of the dose resulting from exposure to site media to the dose believed to be safe, even in sensitive individuals i.e. children and elderly. Here, the  $HRI < 1$  shows that no significant risk of chronic-toxic effects exists, while, the  $HRI > 1$  shows that chronic-toxic effects may occur (18, 19).

In view of the concern for food safety, this study was conducted to analyze the Cd, Cu, Pb, and Zn

contents in raw and pasteurized cows' milk marketed Hamadan City, western Iran, in 2014.

## **Materials and Methods**

### *Reagents*

Standard stock solutions of analyzed metal ions at the concentration of 1000 ppm were used to prepare working solutions after appropriate dilution. Standard solutions were of analytical grade (Merck, Darmstadt, Germany). Distilled deionized water was used in all dilution procedures

### *Sampling and Sample Analysis*

In this study, a total of 36 samples of raw (from the 12 dairies) and 36 samples of pasteurized (from 12 brands) cow's milk were collected from market basket of Hamadan City, western Iran in 2014. Milk samples were collected in sterile polypropylene bottles of 50 ml and they were frozen under  $-20\text{ }^{\circ}\text{C}$  until their analysis (1).

In the laboratory, to obtain the total dehydration of samples, all milk samples were lyophilized. Moreover, for the acid digestion of samples, a microwave system was used. Then 10.0 ml of  $\text{HNO}_3$  (65%) was added to a dry sample of 0.5 gr of the lyophilized material. The solution was kept in the microwave for two periods of 5 min and 10 min to  $100\text{ }^{\circ}\text{C}$  and  $175\text{ }^{\circ}\text{C}$ , respectively with an intermediate depressurization. The solution was kept to  $4\text{ }^{\circ}\text{C}$  up to the moment of its valuation (1, 3). The analysis of Cd, Cu, Pb, and Zn were performed by ICP-OES (Varian, 710-ES) in  $\mu\text{g}/\text{kg}$ . To control the accuracy of the experiment, a multi-element standard solution (Merck) with different concentrations of Cd, Cu, Pb and Zn (0.2, 1, 10, 50, 100 and  $500\text{ }\mu\text{g}/\text{L}$ ) were used for the calibration. For the half of the measurement, three controls of each metal are used, a blank and a standard of  $10\text{ }\mu\text{g}/\text{L}$  (similar to the calibration) (1).

### *Human Health Risk Assessment*

For human health risk assessment the average daily intake of metal (DIM) was calculated using the equation 1 (20-22):

$$DIM = \frac{C_{\text{metal}} \times C_{\text{factor}} \times D_{\text{food intake}}}{B_{\text{average weight}}} \quad (1)$$

, where  $C_{\text{metal}}$ ,  $C_{\text{factor}}$ ,  $D_{\text{food intake}}$ , and  $B_{\text{average weight}}$  are the heavy metal contents in milk ( $\mu\text{g}/\text{kg}$ ), conversion factor (0.085) (21), average daily intake of milk (0.14 kg per person per day) (23), and average body weight (15.0 kg for children and 70.0 kg for adult), respectively (24-26).

The health risk index (HRI) for the population of study area through the consumption of cow's milk was assessed in accordance with equation 2 (20-22):

$$HRI = \frac{DIM}{RfD} \quad (2)$$

, where DIM and RfD are daily intake of metal and oral reference dose of metal, respectively. The RfD for Cd, Cu, Pb, and Zn were 1.0, 40.0, 3.50, and 330.0  $\mu\text{g}/\text{kg}/\text{day}$ , respectively. Here, if the  $HRI < 1$ , is assumed the exposed populations to be safe (19, 22, 26).

The total HRI (THRI) of heavy metals for the milk was calculated according to equation 3 (27):

$$THRI = HRI (\text{toxicant } 1) + HRI (\text{toxicant } 3) + \dots + HRI (\text{toxicant } n) \quad (3)$$

### Statistical Analyses

The statistical analysis of the obtained results consisted in a first Shapiro-Wilk test for normality, followed by the study of the one-way ANOVA to examine statistical significance of differences in the mean concentration of analyzed metals in milk samples. Moreover, concentration of the analyzed metals in cow's milk between raw and pasteurized samples was compared by the independent t-test. Finally, to study a correlation between the metals in the different cow's milk samples, 2-tailed test of Pearson correlation was performed.

### Results

The contents of Cd, Cu, Pb and Zn in the milk samples are presented in Table 1.

**Table 1:** Heavy metal concentrations (mean  $\pm$  S.D.) of the milk samples ( $\mu\text{g}/\text{kg}$ , DW)

Sample	Metal Concentration			
	Cd	Cu	Pb	Zn
		<i>Raw Milk</i>		
1	0.78 $\pm$ 0.01 <sup>e</sup>	15.50 $\pm$ 0.50 <sup>e</sup>	66.90 $\pm$ 0.43 <sup>e</sup>	364.0 $\pm$ 0.00 <sup>e</sup>
2	0.78 $\pm$ 0.01 <sup>e</sup>	15.90 $\pm$ 0.72 <sup>e</sup>	67.30 $\pm$ 0.64 <sup>e</sup>	364.0 $\pm$ 0.58 <sup>e</sup>
3	0.78 $\pm$ 0.01 <sup>e</sup>	16.60 $\pm$ 0.06 <sup>f</sup>	68.00 $\pm$ 0.00 <sup>f</sup>	364.0 $\pm$ 0.00 <sup>e</sup>
4	0.45 $\pm$ 0.02 <sup>d</sup>	8.00 $\pm$ 0.00 <sup>b</sup>	28.00 $\pm$ 0.15 <sup>d</sup>	278.0 $\pm$ 0.00 <sup>d</sup>
5	0.42 $\pm$ 0.01 <sup>c</sup>	8.00 $\pm$ 0.06 <sup>b</sup>	28.60 $\pm$ 0.02 <sup>d</sup>	278.0 $\pm$ 0.58 <sup>d</sup>
6	0.44 $\pm$ 0.01 <sup>d</sup>	8.00 $\pm$ 0.00 <sup>b</sup>	29.00 $\pm$ 0.00 <sup>d</sup>	278.0 $\pm$ 0.00 <sup>d</sup>
7	0.06 $\pm$ 0.01 <sup>a</sup>	9.20 $\pm$ 0.21 <sup>c</sup>	19.00 $\pm$ 0.02 <sup>b</sup>	252.0 $\pm$ 1.00 <sup>e</sup>
8	0.06 $\pm$ 0.01 <sup>a</sup>	9.10 $\pm$ 0.61 <sup>c</sup>	19.20 $\pm$ 0.55 <sup>b</sup>	249.0 $\pm$ 4.73 <sup>b</sup>
9	0.06 $\pm$ 0.01 <sup>a</sup>	10.00 $\pm$ 0.02 <sup>d</sup>	19.80 $\pm$ 0.26 <sup>c</sup>	251.7 $\pm$ 0.58 <sup>bc</sup>
10	0.18 $\pm$ 0.01 <sup>b</sup>	5.70 $\pm$ 0.00 <sup>a</sup>	15.70 $\pm$ 0.06 <sup>a</sup>	122.0 $\pm$ 0.00 <sup>a</sup>
11	0.18 $\pm$ 0.01 <sup>b</sup>	5.70 $\pm$ 0.06 <sup>a</sup>	15.70 $\pm$ 0.17 <sup>a</sup>	122.0 $\pm$ 0.00 <sup>a</sup>
12	0.18 $\pm$ 0.00 <sup>b</sup>	5.70 $\pm$ 0.06 <sup>a</sup>	15.90 $\pm$ 0.06 <sup>a</sup>	122.0 $\pm$ 0.00 <sup>a</sup>
<b>Mean Concentration<math>\pm</math>S.D.</b>	0.36 $\pm$ 0.28	9.77 $\pm$ 3.91	32.83 $\pm$ 20.80	253.70 $\pm$ 87.96
		<i>Pasteurized Milk</i>		
1	0.01 $\pm$ 0.00 <sup>a</sup>	5.10 $\pm$ 0.00 <sup>bc</sup>	1.10 $\pm$ 0.00 <sup>b</sup>	60.10 $\pm$ 0.00 <sup>d</sup>
2	0.01 $\pm$ 0.00 <sup>a</sup>	5.10 $\pm$ 0.00 <sup>bc</sup>	1.10 $\pm$ 0.00 <sup>b</sup>	60.00 $\pm$ 0.05 <sup>d</sup>
3	0.01 $\pm$ 0.00 <sup>a</sup>	5.10 $\pm$ 0.00 <sup>bc</sup>	1.10 $\pm$ 0.03 <sup>c</sup>	61.00 $\pm$ 0.00 <sup>e</sup>
4	0.20 $\pm$ 0.00 <sup>a</sup>	5.00 $\pm$ 0.00 <sup>abc</sup>	1.00 $\pm$ 0.00 <sup>a</sup>	30.20 $\pm$ 0.29 <sup>b</sup>
5	0.20 $\pm$ 0.00 <sup>a</sup>	5.00 $\pm$ 0.00 <sup>abc</sup>	1.00 $\pm$ 0.00 <sup>a</sup>	30.20 $\pm$ 0.29 <sup>b</sup>
6	0.14 $\pm$ 0.10 <sup>a</sup>	5.00 $\pm$ 0.06 <sup>bc</sup>	1.10 $\pm$ 0.00 <sup>b</sup>	31.00 $\pm$ 0.00 <sup>c</sup>
7	21.20 $\pm$ 0.29 <sup>c</sup>	18.50 $\pm$ 0.10 <sup>d</sup>	63.0 $\pm$ 0.00 <sup>e</sup>	25.00 $\pm$ 0.00 <sup>a</sup>
8	21.30 $\pm$ 0.58 <sup>c</sup>	18.40 $\pm$ 0.26 <sup>d</sup>	63.0 $\pm$ 0.00 <sup>e</sup>	25.00 $\pm$ 0.00 <sup>a</sup>
9	22.00 $\pm$ 0.00 <sup>d</sup>	18.90 $\pm$ 0.10 <sup>d</sup>	63.0 $\pm$ 0.00 <sup>e</sup>	25.00 $\pm$ 0.00 <sup>a</sup>
10	0.60 $\pm$ 0.00 <sup>b</sup>	4.30 $\pm$ 0.58 <sup>a</sup>	37.0 $\pm$ 0.00 <sup>d</sup>	244.70 $\pm$ 0.57 <sup>g</sup>
11	0.59 $\pm$ 0.00 <sup>b</sup>	4.70 $\pm$ 1.15 <sup>ab</sup>	37.0 $\pm$ 0.00 <sup>d</sup>	244.30 $\pm$ 0.57 <sup>f</sup>
12	0.59 $\pm$ 0.00 <sup>b</sup>	5.70 $\pm$ 0.10 <sup>c</sup>	37.0 $\pm$ 0.00 <sup>d</sup>	245.00 $\pm$ 0.00 <sup>g</sup>
<b>Mean Concentration<math>\pm</math>S.D.</b>	5.57 $\pm$ 9.33	8.41 $\pm$ 5.99	25.54 $\pm$ 26.50	90.12 $\pm$ 91.52

\* The letters (a, b, c, d ...) represent the significant difference between the mean concentration of metals in cow's milk samples that computed by One-way ANOVA and Duncan multiple range test ( $p = 0.05$ )

Based on the results, among the raw cow's milk samples, Cd, Cu, Pb and Zn ( $\mu\text{g}/\text{kg}$ ) were detected in amount ranging from 0.06 to 0.78, 5.70 to 16.60, 15.70 to 68.0, and 122.0 to 364.0, respectively. In addition, among the pasteurized cow's milk samples, Cd, Cu, Pb and Zn ( $\mu\text{g}/\text{kg}$ ) were detected an amount ranging from 0.01 to 22.0, 4.30 to 18.90, 1.0 to 63.0, and 25.0 to 245.0, respectively.

Comparing the heavy metal contents in milk samples with the maximum permissible limits (MPL) (2.6  $\mu\text{g}/\text{kg}$  for Cd, 10  $\mu\text{g}/\text{kg}$  for Cu, 20  $\mu\text{g}/\text{kg}$  for Pb and 328  $\mu\text{g}/\text{kg}$  for Zn, respectively.) established by International Dairy Federation (28), indicated that the mean concentration of Cd in raw cow's milk samples and Cu and Zn in raw and pasteurized cow's milk samples, were lower than MPL.

In addition, HRI values in adults and children through consumption of both raw and pasteurized cow's milk were lower than 1 (safe limits) (Table 2). Furthermore, the total HRI values (THRI) of heavy metals for adults via consumption of raw cow's milk, which varied from 8.66E-04 to 3.71E-03 and for children were varied from 4.04E-03 to 1.73E-02 and below the safe limit (THRI < 1). Besides, the THRI values for adults through consumption of pasteurized cow's milk, which varied from 8.28E-05 to 7.02E-03 and for children, were varied from 1.21E-02 to 3.27E-02 and similar to raw milk samples below the safe limit (THRI < 1). Therefore, all consumers including adults and children have no potential health risk through consuming raw and pasteurized cow's milk from the study area.

**Table 2:** Daily intakes of metals (DIM,  $\mu\text{g}$ ) and health risk index (HRI) for individual heavy metal caused by the raw and pasteurized cow's milk

	<i>Cd</i>	<i>Cu</i>	<i>Pb</i>	<i>Zn</i>
<i>Raw Milk</i>				
Adults				
DIM	6.12E-05	1.66E-03	5.58E-03	3.10E-01
STD	4.76E-05	6.65E-04	3.54E-03	1.50E-02
Min	1.02E-05	9.69E-04	2.67E-03	2.10E-02
Max	1.33E-04	2.82E-03	1.16E-02	6.20E-02
HRI	6.12E-05	4.15E-05	1.59E-03	1.03E-03
STD	4.76E-05	1.66E-05	1.01E-03	4.98E-05
Min	1.02E-05	2.42E-05	7.62E-04	6.91E-05
Max	1.33E-04	7.05E-05	3.30E-03	2.10E-04
Children				
DIM	2.86E-04	7.75E-03	2.60E-02	2.01E-01
STD	2.22E-04	3.10E-03	1.65E-02	6.98E-02
Min	4.76E-05	4.52E-03	1.24E-02	9.68E-02
Max	6.20E-04	1.32E-02	5.39E-02	2.89E-01
HRI	2.86E-04	1.94E-04	7.44E-03	6.71E-04
STD	2.22E-04	7.75E-05	4.71E-03	2.33E-04
Min	4.76E-05	1.13E-04	3.56E-03	3.22E-04
Max	6.20E-04	3.29E-04	1.54E-02	9.62E-04
<i>Pasteurized Milk</i>				
Adults				
DIM	9.47E-04	1.43E-03	4.34E-03	1.53E-02
STD	1.59E-03	1.02E-03	4.51E-03	1.55E-02
Min	1.70E-06	7.31E-04	1.70E-04	4.25E-04
Max	3.74E-03	3.21E-03	1.07E-04	4.16E-02
HRI	9.46E-04	3.57E-05	1.24E-03	5.10E-05
STD	1.59E-03	2.54E-05	1.29E-03	5.19E-05
Min	1.70E-06	1.83E-05	4.86E-05	1.42E-05
Max	3.74E-03	8.03E-05	3.06E-03	1.39E-04
Children				
DIM	4.42E-03	6.67E-03	2.03E-02	7.15E-02
STD	7.40E-03	4.75E-03	2.10E-02	7.26E-02
Min	7.93E-06	4.72E-01	7.93E-04	1.98E-02
Max	1.74E-02	1.50E-02	5.00E-02	1.94E-01
HRI	4.42E-03	1.67E-04	5.79E-03	2.38E-04
STD	7.40E-03	1.19E-04	6.00E-03	2.42E-04
Min	7.93E-06	1.18E-02	2.27E-04	6.61E-05
Max	1.74E-02	3.75E-04	1.43E-02	6.47E-04

Based on the results of the Pearson's correlations coefficient in the raw and pasteurized milk samples, there are significantly correlated were found between concentrations of some metals.

The results of independent t-test showed that there are significantly different ( $P < 0.05$ ) were found in the contents of Cd and Zn between the raw and pasteurized cow's milk samples. While, for the levels of Cu and Pb no significant differences ( $P > 0.05$ ) between the raw and pasteurized cow's milk samples were found.

## Discussion

In this study, the 36 raw and 36 pasteurized milk samples were analyzed for determination of Cd, Cu, Pb and Zn content after the microwave acid extraction. Cadmium is very toxic metal with a natural occurrence or human activities origin in the environment (29). Acute toxicities of Cd cause deflection of cardiovascular and skeleton

systems (8). The mean concentrations of Cd in raw and pasteurized cow's milk samples ( $\mu\text{g}/\text{kg}$ ) were  $0.36 \pm 0.28$  and  $5.57 \pm 9.33$ , respectively and that was higher than the recommended standards were found in pasteurized cow's milk samples. Cadmium accumulation in plants as the animal feed grown near industrial areas, zones of storage of mining waste depots and regions with high traffic volume can translocation the Cd in the animals' tissues, and consequently in the milk (1, 30, 31). Moreover, except for raw materials, the equipment used during the production process and packaging materials are the possible sources of contamination (5). The obtained results are similar to the findings that concluded the concentrations of Cd in raw milk samples collected from Serbia ( $3.50 \mu\text{g}/\text{kg}$ ) were much lower than Serbian regulations maximal permitted contents ( $10 \mu\text{g}/\text{kg}$ ) (5). Comparison of our findings with other studies is shown in Table 3.

**Table 3:** The metal contents ( $\mu\text{g}/\text{kg}$ ) in raw and pasteurized milk samples compared to other studies

Region	Elements				References
	Cd	Cu	Pb	Zn	
Iran (Raw milk)	0.36	9.77	32.83	253.70	Present study
Iran (Pasteurized Milk)	5.57	8.41	25.54	90.12	Present study
Egypt	246	1448	3141	7203	(28)
Poland (Simmental cow)	3.50	37.70	36.60	3027	(42)
Poland (Holstein-Friesian cow)	4.00	45.30	41.20	3277	(42)
Spain	<2	-	3.91	-	(1)
South Africa	<0.006	-	5.44	-	(32)
Lithuania (winter season)	0.37	-	0.47	-	(33)
Lithuania (Summer season)	0.18	-	0.54	-	(33)
Croatian	5.31	-	34.00	-	(3)
Iran (Hamadan Province)	3.21	-	4.84	-	(34)
Iran (Raw Milk)	2.87	-	60.72	-	(35)
Iran (Pasteurized Milk)	1.03	-	13.57	-	(35)
Iran (Raw Milk)	104.00	142.00	2720.00	3072.00	(36)
China	305.00	-	-	-	(37)
Pakistan (Area 1)	89.00	-	21781	-	(38)
Pakistan (Area 2)	62.00	-	15958	-	(38)
Denmark	0.66	-	-	-	(39)

Copper known as essential micronutrients for growing plants should be ensured through organic and artificial fertilizers or which is important to healthy hormone secretion, nerve conduction, and the growth of bones and connective tissue. As the literature shows, Cu plays a critical role in some biochemical processes (9, 10). Based on the

results, the concentrations of Cu in raw and pasteurized cow's milk samples ( $\mu\text{g}/\text{kg}$ ) with an average of  $9.77 \pm 3.91 \mu\text{g}/\text{kg}$  and  $8.41 \pm 5.99 \mu\text{g}/\text{kg}$  were lower than the MPL. This reported of Cu level was much lower than those observed in raw milk samples collected from Serbia ( $117.70 \mu\text{g}/\text{kg}$ ) (5). Although, the mean concentrations of

Cu were below the MPL, animal feed, drinking water with the high amount of Cu and Cu alloys used in equipment is a possible source of contamination of Cu in milk. (28). Comparison of our findings with other studies is shown in Table 3.

Lead as a highly toxic metal has no known functions in biological systems and recognized as a major environmental health risk throughout the world (40). The concentrations of Pb in raw and pasteurized cow's milk samples ( $\mu\text{g}/\text{kg}$ ) with an average of  $32.83 \pm 20.80$  and  $25.54 \pm 26.50$ , respectively were higher than the recommended standards. Similarly, like for Cd, the highest Pb levels in milk samples were reported in animals reared in the vicinity of Pb–Zn smelter, mining waste depots, thermal power and roads usually exposed to more automotive Pb emissions (1). In this regard, Suturovic et al. (2014) reported that the Pb contents in raw milk samples were  $75.42 \mu\text{g}/\text{kg}$ . Comparison of our findings with other studies is shown in Table 3.

Zinc is known as vital element for human growth (14). The concentrations of Zn in raw and pasteurized cow's milk samples ( $\mu\text{g}/\text{kg}$ ) with an average of  $253.70 \pm 87.96$  and  $90.12 \pm 91.52$ , respectively were lower than the MPL. These values were lower than findings that reported the variation in Zn content in milk samples collected from Egypt ranged from  $888 \mu\text{g}/\text{kg}$  to  $18316 \mu\text{g}/\text{kg}$  (41). Comparison of our findings with other studies is shown in Table 3.

The average HRI values through consumption of raw cow's milk were  $6.81\text{E-}04$  and  $2.15\text{E-}03$  for adults and children, respectively (Table 2). While, the average HRI values through consumption of pasteurized cow's milk were  $5.68\text{E-}04$  and  $2.65\text{E-}03$  for adults and children, respectively. Therefore, mean of HRI values of analyzed metals for adults and children through consumption of raw and pasteurized cow's milk are lower than 1. In this regard, the HRI of all analyzed metals were minimal.

The Pearson's correlations coefficient were performed between metal concentrations in raw and pasteurized cow's samples to understand the relationships between them. In the raw milk samples the residual content of Cd was significantly positively correlated with the contents of Cu

( $r=0.779$ ), Pb ( $r=0.939$ ) and Zn ( $r=0.760$ ), the residual content of Cu was significantly positively correlated with the concentrations of Pb ( $r=0.941$ ) and Zn ( $r=0.884$ ) and the residual content of Pb was significantly positively correlated with the concentration of Zn ( $r=0.842$ ). Moreover, in the pasteurized milk samples the residual content of Cd was significantly positively correlated with the concentrations of Cu ( $r=0.997$ ) and Pb ( $r=0.840$ ) and negatively correlated with the concentration of Zn ( $r=-0.397$ ), and the residual content of Cu was significantly positively correlated with the concentration of Pb ( $r=0.819$ ) and the negative correlated with the concentration of Zn ( $r=-0.425$ ), proved in our work. In this regard, strong positive correlations were reported between Pb and Cd concentrations ( $r=0.85$  vs.  $r=0.87$ ) in the milk of Simmental and Holstein-Friesian breeds (42). The residual content of Cu in milk was significantly positively correlated with the concentrations of Zn ( $r=0.629$ ) (43). However, significant negative correlations were found between Cu and Zn concentrations ( $r=-0.377$ ) (44).

## Conclusion

This study was carried out to analyze the Cd, Cu, Pb, and Zn contents in raw and pasteurized cow's milk from city of Hamadan, Iran. Based on the results, although in the pasteurized cow's milk samples, the contents of Cu, Pb, and Zn were lower compared to the raw cow's milk samples, but considering the serious contamination of three samples of pasteurized milk by Cd, four samples of raw milk and three samples of pasteurized milk by Cu, six samples of both raw and pasteurized milk by Pb and three samples of raw milk by Zn (above the MPL) a control of heavy metals content during the whole production process of milk must be applied. In addition, to maintain human health, it is recommended that special attention be given to the adverse effect of heavy metals through the consumption of other foodstuffs.

## Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or

falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the author.

## Acknowledgements

The author is grateful to the Hamadan Branch, Islamic Azad University for providing facilities to conduct and complete this study.

## Conflict of interest

The author declares that there is no conflict of interest.

## References

1. González-Montaña JR, Senís E, Gutiérrez A, Prieto F (2012). Cadmium and lead in bovine milk in the mining area of the Caudal River (Spain). *Environ Monit Assess*, 184(7): 4029-4034.
2. Hosseini SV, Sobhanardakani S, Tahergorabi R, Delfieh P (2013). Selected heavy metals analysis of Persian sturgeon's (*Acipenser persicus*) caviar from Southern Caspian Sea. *Biol Trace Elem Res*, 154(3): 357-362.
3. Pavlovic I, Sikiric M, Havranek JL et al (2004). Lead and cadmium levels in raw cow's milk from an industrialised Croatian region determined by electrothermal atomic absorption spectrometry. *Czech J Anim Sci*, 49(4): 164-168.
4. Swarup D, Patra RC, Naresh R et al (2005). Blood lead levels in lactating cows reared around polluted localities; transfer of lead into milk. *Sci Total Environ*, 347(1-3):106-10.
5. Suturovic Z, Kravic S, Milanovic S et al (2014). Determination of heavy metals in milk and fermented milk products by potentiometric stripping analysis with constant inverse current in the analytical step. *Food Chem*, 155: 120-125.
6. Licata P, Trombetta D, Cristani M et al (2004). Levels of "toxic" and "essential" metals in samples of bovine milk from various dairy farms in Calabria, Italy. *Environ Int*, 30(1): 1-6.
7. Ghindini S, Zanardi E, Battaglia A et al (2005). Comparison of contaminant and residue levels in organic and conventional milk and meat products from Northern Italy. *Food Addit Contam*, 22(1): 9-14.
8. Özcan MM, AL Juhaimi FY (2012). Determination of heavy metals in bee honey with connected and not connected metal wires using inductively coupled plasma atomic emission spectrometry (ICP-AES). *Environ Monit Assess*, 184(4): 2373-2375.
9. Saracoglu S, Tuzen M, Soylak M (2009). Evaluation of trace element contents of dried apricot samples from Turkey. *J Hazard Mater*, 167(1-3): 647-652.
10. Ackah M, Anim AK, Zakaria N et al (2014). Determination of some heavy metal levels in soft drinks on the Ghanaian market using atomic absorption spectrometry method. *Environ Monit Assess*, 186(12): 8499-8507.
11. Needleman H (2004). Lead poisoning. *Annu Rev Med*, 55: 209-222.
12. Finster ME, Kimberly AG, Helen JB (2004). Lead levels of edibles grown in contaminated residential soils: a field survey. *Sci Total Environ*, 320(2-3): 245-257.
13. Liu WT, Zhou QX, Zhang YL, Wei S (2010). Lead accumulation in different Chinese cabbage cultivars and screening for pollution-safe cultivars. *J Environ Manage*, 91(3): 781-788.
14. Grattan BJ, Freake HC (2012). Zinc and cancer: Implications for LIV-1 in breast cancer. *Nutrients*, 4(7): 648-675.
15. Sobhanardakani S, Jamshidi K (2015). Assessment of metals (Co, Ni, and Zn) content in the sediments of Mighan Wetland using geoaccumulation index. *Iran J Toxicol*, 9(30):1386-1390.
16. World Health Organization (2010). WHO Human Health Risk Assessment Toolkit: Chemical Hazards. International Programme on Chemical Safety. [www.who.int/ipcs/methods/harmonization/areas/ra\\_toolkit/en/](http://www.who.int/ipcs/methods/harmonization/areas/ra_toolkit/en/)
17. Sobhanardakani S (2016). Health risk assessment of As and Zn in canola and soybean oils consumed in Kermanshah City. *J Adv Environ Health Res*, 4(2): 62-67.
18. Ru QM, Feng Q, He JZ (2013). Risk assessment of heavy metals in honey consumed in Zhejiang province, southeastern China. *Food Chem Toxicol*, 53: 256-262.

19. Liang Q, Xue ZJ, Wang F et al (2015). Contamination and health risks from heavy metals in cultivated soil in Zhangjiakou City of Hebei Province, China. *Environ Monit Assess*, 187(12): 754.
20. Khan S, Cao Q, Zheng Y, Huang YZ, Zhu YG (2008). Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China. *Environ Pollut*, 152(3): 686-692.
21. Jan FA, Ishaq M, Khan S et al (2010). A comparative study of human health risks via consumption of food crops grown on wastewater irrigated soil (Peshawar) and relatively clean water irrigated soil (lower Dir). *J Hazard Mater*, 179(1-3): 612-621.
22. Guo J, Yue T, Li X, Yuan Y (2016). Heavy metal levels in kiwifruit orchard soils and trees and its potential health risk assessment in Shaanxi, China. *Environ Sci Pollut Res Int*, 23(14): 14560-6.
23. Abdi F, Atarodi Kashani Z, Mirmiran P, Esteki T (2015). Surveying global and Iranian food consumption patterns: A review of the literature. *J Fasa Univ Med Sci*, 5(2): 159-167. [Persian]
24. Falco G, Ilobet J, Bocio A, Domingo JL (2006). Daily intake of arsenic, cadmium, mercury, and lead by consumption of edible marine species. *J Agric Food Chem*, 54(16):6106-12.
25. Omar WA, Zaghoul KH, Abdel-Khalek AA, Abo-Hegab S (2013). Risk assessment and toxic effects of metal pollution in two cultured and wild fish species from highly degraded aquatic habitats. *Arch Environ Contam Toxicol*, 65(4): 753-764.
26. Tang W, Cheng J, Zhao W, Wang W (2015). Mercury levels and estimated total daily intakes for children and adults from an electronic waste recycling area in Taizhou, China: Key role of rice and fish consumption. *J Environ Sci (China)*, 34:107-15.
27. Saha N, Zaman M (2013). Evaluation of possible health risks of heavy metals by consumption of foodstuffs available in the central market of Rajshahi City, Bangladesh. *Environ Monit Assess*, 185(5): 3867-3878.
28. Malhat F, Hagag M, Saber A, Fayz AE (2012). Contamination of Cows milk by heavy metal in Egypt. *Bull Environ Contam Toxicol*, 88(4): 611-613.
29. Zhu F, Fan W, Wang X et al (2011). Health risk assessment of eight heavy metals in nine varieties of edible vegetable oils consumed in China. *Food Chem Toxicol*, 49(12): 3081-3085.
30. Parkpian P, Leong ST, Laortanakul P, Thunthaisong N (2003). Regional monitoring of lead and cadmium contamination in a tropical grazing land site, Thailand. *Environ Monit Assess*, 85(2): 157-173.
31. Simsek O, Gultekin R, Oksuz O, Kurultay S (2000). The effect of environmental pollution on the heavy metal content of raw milk. *Nahrung*, 44(5): 360-363.
32. Ataro A, McCrindle RI, Botha BM et al (2008). Quantification of trace elements in raw cow's milk by inductively coupled plasma mass spectrometry (ICP-MS). *Food Chem*, 111(1): 243-248.
33. Valiukenaite R, Stankeviciene M, Stankevicius H, Skibniewska KA (2006). Lead and cadmium were determined by inductively coupled plasma sector field mass spectrometry. *Pol J Food Nutr Sci*, 56: 243-246.
34. Vahidinia AA, Salehi I, Beyginagad H et al (2013). Assessment of lead and cadmium contamination and influencing factors in raw milk from regions of Hamadan province. *J Food Hyg*, 3(3): 39-47. [Persian]
35. Bonyadian M, Moshtaghi H, Soltany Z (2006). Study on the residual of lead and cadmium in raw and pasteurized milks in Shahrekord area. *Iran J Vet Res*, 2(2): 74-81. [Persian]
36. Sobhanardakani S, Tizhosh M (2016). Determination of Zn, Pb, Cd and Cu contents in raw milk from the Khorram Abad dairies. *J Food Hyg*, 6(2): 43-50. [Persian]
37. Sun WC, Luo YH, Ma HQ (2011). Preliminary study of metal in yak (*Bos grunniens*) milk from Qilian of the Qinghai Plateau. *Bull Environ Contam Toxicol*, 86(6): 653-656.
38. Javed I, Jan I, Muhammad F et al (2009). Heavy metal residues in the milk of cattle and goats during winter season. *Bull Environ Contam Toxicol*, 82(5): 616-620.
39. Hermansen JE, Badsberg JH, Kristensen T, Gundersen V (2005). Major and trace elements in organically or conventionally produced milk. *J Dairy Res*, 72(3): 362-368.
40. Matovic V, Buha A, Đukić-Ćosić D, Bulat Z (2015). Insight into the oxidative stress induced by lead and/or cadmium in blood,



- liver and kidneys. *Food Chem Toxicol*, 78: 130-140.
41. Meshref AMS, Moselhy WA, El-Houda N, Hassan Y (2014). Heavy metals and trace elements levels in milk and milk products. *J Food Meas Charact*, 8(4): 381-388.
  42. Pilarczyk R, Wójcik J, Czerniak P et al (2013). Concentrations of toxic heavy metals and trace elements in raw milk of Simmental and Holstein-Friesian cows from organic farm. *Environ Monit Assess*, 185(10): 8383-8392.
  43. Sikirić M, Brajenović N, Pavlović I et al (2003). Determination of metals in cow's milk by flame atomic absorption spectrophotometry. *Czech J Anim Sci*, 48(11): 481-486.
  44. Rodríguez Rodríguez EM, Sanz Alaejos M, Díaz Romero C (1999). Chemometric studies of several minerals in milks. *J Agric Food Chem*, 47(4): 1520-4.