



Effects of Hair Metals on Body Weight in Iranian Children Aged 20 to 36 Months

*Mohsen VIGEH^{1,2,3}, Kazuhito YOKOYAMA², Takehisa MATSUKAWA², Atsuko SHINOHARA², Mamak SHARIAT¹, Katsumi OHTANI³

1. Maternal, Fetal & Neonatal Research Center, Imam Khomeini Hospital, Tehran University of Medical Sciences, Tehran, Iran
2. Dept. of Epidemiology and Environmental Health, Juntendo University Faculty of Medicine, 2-1-1 Hongo, Bunkyo-ku, Tokyo, Japan
3. Occupational Epidemiology Research Group, National Institute of Occupational Safety and Health, 6-21-6 Nagao, Tama-ku, Kawasaki, Japan

*Corresponding Author: Email: mohsen.vigeh@gmail.com

(Received 11 Jul 2016; accepted 10 Dec 2016)

Abstract

Background: Although the level of exposure to many toxic metals decreased recently, the adverse effects of these metals on children's growth and development remain a serious public health issue.

Methods: The present study was conducted in three teaching hospitals affiliated with Tehran University of Medical Sciences (Tehran, Iran) from Sep 2012 to Mar 2013. To study the relationship between metals and childhood growth, concentrations of zinc and several potentially toxic metals (lead, cadmium, antimony, cobalt, and molybdenum) were measured in scalp hair for 174 children, aged 20 to 36 months.

Results: The hair concentrations of cobalt were significantly ($P < 0.05$) higher in children at the lower percentile of weight than in higher-weight children (0.026 ± 0.04 vs. 0.015 ± 0.01 $\mu\text{g/g}$, respectively). Hair contents of lead, cobalt, and antimony were significantly higher ($P < 0.05$) in girls than in boys (8.08 ± 8.7 vs. 4.92 ± 5.6 $\mu\text{g/g}$ for lead, 0.026 ± 0.03 vs. 0.16 ± 0.02 $\mu\text{g/g}$ for cobalt, and 0.188 ± 0.29 vs. 0.102 ± 0.12 $\mu\text{g/g}$ for antimony). There were also significant correlations between lead and other metals in the children's hair.

Conclusion: Gender may play a significant role in absorption and/or accumulation of metals. It should be considered when we study metal toxicity in children.

Keywords: Metal, Children, Gender, Growth, Weight, Hair

Introduction

Childhood exposure to toxic metals is a serious public health problem in many developing countries (1) and usually, involves exposure to a mixture rather than individual metals (2). Thus, it is critical to investigate metals interactions due to mixed metals exposure has been reported to increase the risk of adverse effects on children's growth and development (3). However, few epidemiological surveys have studied multiple metals exposure and their interaction in children.

Personal characteristics, such as age and sex, have been reported to influence metals' metabolisms, which may induce metal toxicities (4-9).

This may occur via synergism in the production of reactive oxygen species, cytokine pathway activation, and cytochrome P450 enzyme activity (5). For instance, glutathione (an antioxidant agent) plays a central role against biotic stress in coping with different metals (6, 7), especially when a multiple metal exposure occurs (8). Epidemiological (9, 10) and experimental (rats) (11) studies have shown significant differences in levels of glutathione and superoxide dismutase between female and male subjects, which may lead to differing in protection against oxidative stress (12).

Studies on preschool children found considerable numbers of underweight children (7.5% to 20%) (13-16) and children of short stature (6.6%) (13) in Iran compared to international data (17), with a significant difference between boys and girls (16). The reasons for Iranian children growth problem is not fully understood yet. Although socioeconomic status and subsequent diets insufficiency may play an important role, other factors, such as genes, gender, and environmental pollution, may be involved as well. Iranian infants are exposed to relatively high levels of metals from industrial activities, via polluted air and soils (18, 19) or through breast milk (mean lead level of 2.4 µg/dL) (20). In addition, a recent study in Iran reported higher levels of metal in the first haircut of newborn than the mothers' hair (157 vs 87 µg/kg for cadmium, 246 vs 198 µg/kg for mercury, 14313 vs 11776 µg/kg for copper, and 408207 vs 52022 µg/kg for aluminum, respectively) (21), which can suggest a free pass of these metals via umbilical cord to the fetuses.

Serial blood measurements may offer a better estimation of the body burden of metals than a single measurement, but they are usually impractical. Instead, measuring metals in hair samples offers advantages for routine clinical screening and diagnosis as an alternative to blood and urine sampling (22-24), which can present information on exposure over a long period of time and non-invasive sample collection.

The present study measured concentrations of metals (lead, zinc, cadmium, antimony, cobalt, and molybdenum) in children's scalp hair to evaluate metal effects on children's weight gain.

Materials and Methods

Study participants

The present study was conducted in three teaching hospitals affiliated with Tehran University of Medical Sciences (Tehran, Iran) from Sep 2012 to Mar 2013. The study included 174 children aged 20 months to 3 yr.

The Ethical Committees for the Vice-Chancellor of the Research Department and Institutional Review Board of Tehran University of Medical

Sciences approved the study design and procedures. The study was conducted under the supervision of the Maternal, Fetal & Neonatal Research Center, Tehran University of Medical Sciences. The study purpose and procedures were fully explained to the children's parents, and the study was conducted with their informed consent. Participation in the study was strictly voluntary.

Questionnaire and measurements

Information about participants' characteristics was gathered using a structured questionnaire developed for this study (see Appendix for main items). Anthropometric variables were obtained from growth chart records. Children were weighed when wearing underwear only, using a standard balance beam scale. With the children in standing position, height was measured to the nearest 0.1 cm using a rigid stadiometer tested for accuracy.

Collection and analysis of hair samples

For removing external contamination from children's hair, mothers were asked to wash the children's hair with shampoo or soap, then rinse with plenty of tap water (at least five minutes) immediately prior to sampling. None used any type of hair treatment. After cutting children's hair (about 3 cm from the child's scalp; 2-5 g), the sample was placed in a plastic pack and transferred to the laboratory for metals analysis.

For analysis, hair samples were weighed (5-10 mg) and put into Teflon perfluoroalkoxy bottles, and 0.4 ml of concentrated nitric acid (ultrapure grade, Tama Chemicals Co., Kawasaki, Japan) was added. The bottles were left overnight at room temperature (18-28 °C). The sample mixture was then digested with 0.2 ml of hydrogen peroxide (ultrapure grade, Tama Chemicals Co., Kawasaki, Japan) using a microwave oven (MLS-1200 MEGA, Milestone Srl, Bergamo, Italy). This process was done in five steps using various power levels set at 250, 0, 400, 650, and 250 W for 6, 1, 6, 6, and 6 min, respectively. Next, the volume of the digested sample was adjusted to 1.0 ml using ultrapure water. After dilution with

0.5% nitric acid solution, subsequent measurements for metal concentrations were done by inductively coupled plasma mass spectrometry (ELAN 6000, PerkinElmer, Waltham, MA, USA) using multi-element standard solutions XSTC-13B and XSTC-622 (SPEX CertiPrep, Metuchen, NJ, USA). Each measurement was repeated three times, and the average of the three measurements was used for statistical analyses. For instrument calibration throughout the measurements, at least 10% of the analyses were external standards, and 5% were blank (pure water).

Data analysis

Children weight gain (kg) was calculated by subtracting the children's weight at 18 months old from their birth weight recorded on the growth chart. Pearson correlation coefficients were used to study relationships between metals. Student's t-tests were calculated to investigate differences in metal concentrations between relatively high and low weight gain children (<50th and >50th percentile) at 18 months of age. The same statistical test was employed to compare metal concentrations between boys and girls. We employed

ANOVA test to compare children's hair levels of metals among the city districts. Multiple regression analysis was used to examine the relationships between metal levels and children's weight, controlling for possible confounding variables (maternal age, weight, height, and child's age and sex). Statistical analyses were performed using SPSS (IBM SPSS, Armonk, NY, USA), and statistical significance was determined using $P < 0.05$.

Results

Three children (two boys and one girl) weight were at less than percentile five and nine children (3 boys and 6 girls) weight were higher than percentile 95, according to the World Health Organization (WHO) classification for 18 months old children, separately for boys and girls. Student's t-test showed that children with relatively lower weight at 18 months (<50th percentile) had significantly higher levels of cobalt in their hair than children with higher weight (mean \pm SD 0.026 ± 0.04 and 0.015 ± 0.01 $\mu\text{g/g}$, respectively, $P=0.021$) (Table 1).

Table 1: Comparison of metal levels in children's hair according to percentiles of children weight at 18 months old (n = 138) ‡

Weight percentile	< 50 n= 70†	> 50 n = 68†	P-value
Hair level of ($\mu\text{g/g}$)			
Lead	6.77 ± 6.9	5.14 ± 4.9	NS
Zinc	122 ± 76	123 ± 58	NS
Cadmium	0.110 ± 0.14	$0.087 \pm .07$	NS
Antimony	0.159 ± 0.24	0.110 ± 0.17	NS
Cobalt	0.026 ± 0.04	0.015 ± 0.01	< 0.021
Molybdenum	$0.091 \pm .05$	$0.099 \pm .10$	NS
Child age (month)	27.2 ± 5.0	29.0 ± 4.6	0.031
Maternal			
Age (yr)	25.1 ± 4.6	25.8 ± 4.1	NS
Weight (kg)	61.8 ± 12.8	65.1 ± 9.0	NS
Height (cm)	157.2 ± 7.1	160.2 ± 60.1	0.041

‡.36 missing data (due to unreliable metal measurement according to standards and blank or, missed data on subjects' characteristics)

† Mean \pm SD; Student t-test

NS: none significant

There was not the same result for children's weight percentiles at the birth, 6 and 12 months old. Multiple regression analysis showed no significant relationship between children's weight and hair concentration of cobalt when adjusted for covariates (data not shown). Hair concentrations of lead, cobalt, and antimony were significantly higher in girls than in boys (8.08 ± 8.7 vs. 4.92 ± 5.6 $\mu\text{g/g}$ for lead, 0.026 ± 0.03 vs. $0.16 \pm$

0.02 $\mu\text{g/g}$ for cobalt, and 0.188 ± 0.29 vs. 0.102 ± 0.12 $\mu\text{g/g}$ for antimony, respectively, $P < 0.05$) (Table 2).

There was a significant negative correlation between lead and zinc in children's hair ($r = -0.436$, $P < 0.001$) (Fig. 1). However, lead showed significant positive correlations with potentially toxic metals (cadmium, antimony, cobalt, and molybdenum) in children's hair (Fig. 2).

Table 2: Comparison of metal levels in children's hair, weight, and height between boys and girls (n = 155)[‡]

	Boys n = 74 [†]	Girls n = 81 [†]	P-value
Hair level of ($\mu\text{g/g}$)			
Lead	4.92 ± 5.6	8.08 ± 8.7	0.008
Zinc	124 ± 56	117 ± 72	NS
Cadmium	$.086 \pm .09$	$.118 \pm .12$	NS
Antimony	$.102 \pm .12$	$.188 \pm .29$	0.017
Cobalt	$.016 \pm .02$	$.026 \pm .03$	0.029
Molybdenum	$.088 \pm .05$	$.108 \pm .10$	NS
Child			
Age (month)	26.3 ± 7.1	27.1 ± 5.7	NS
Birth weight (kg)	4.11 ± 4.6	3.43 ± 3.0	NS
Birth height	50.6 ± 2.9	49.2 ± 3.0	0.003
Weight at 6 months (kg)	7.87 ± 0.8	7.3 ± 0.8	0.001
Height at 6 months (cm)	67.2 ± 3.2	65.5 ± 3.5	0.002
Weight at 12 months (kg)	9.8 ± 0.9	9.2 ± 0.9	0.001
Height at 12 months (cm)	75.2 ± 3.4	73.8 ± 3.0	0.002
Weight at 18 months (kg)	11.2 ± 1.1	10.5 ± 1.3	0.001
Height at 18 months (cm)	81.7 ± 3.3	80.1 ± 3.4	0.006
Weight gain (kg)	6.97 ± 4.9	7.04 ± 3.25	NS
Maternal			
Age (year)	25.7 ± 4.0	25.2 ± 4.5	NS
Weight (kg)	62.4 ± 10.5	65.0 ± 11.1	NS
Height (cm)	159.7 ± 6.7	158.1 ± 6.5	NS

[‡]19 missing data (due to unreliable metal measurement according to standards and blank or, missed data on subjects' characteristics) // [†] Mean \pm SD; Student *t*-test // NS: none significant

Among 22 districts of Tehran, the most of study participants live in the district 17 (southwest, n=32) and the district 4 (northeast, n=29). There was not a significant difference between the children scalp hair metals levels or children's weight with the district of living in the city (data not shown).

Discussion

The present study found that relatively lower-weight children had higher levels of cobalt in their hair than children who weighed more at 18

months old. However, this difference disappeared when we adjusted for other variables in multiple regression analysis.

Children's hair levels of some potentially toxic metals were approximately twice as high in girls as boys. Overall, girls also displayed lower weight than boys (Table 2). Although, the present study multivariate statistical analysis failed to demonstrate finding about cobalt and children weight, other study showed the content of metals in body fluids differences between the short stature children and the healthy children (25).




Fig. 1: Pearson correlation between concentrations of lead and zinc in children hair




Fig.2: Pearson correlation between concentrations of lead with metals in children hair

In the other word, gender influencing was reported on metals effect on body weight and/or body mass index (26). For instance, longitudinal studies in Bangladesh have shown that exposure to inorganic arsenic through drinking water during pregnancy and the postnatal period (measured in urine at 18 months) is correlated with decreased size at birth (27) and a lower childhood growth rate (at 18–24 months) (28), and the effect is more pronounced in girls than in boys. Similar findings regarding gender were observed in China, when comparing children based on arsenic levels in their drinking water (2 µg/L arsenic content vs. 190 µg/L) (29). Thus, gender may play an important role in metal toxicokinetics and accumulation, and consequently, in metal toxicities in children (30–32).

As the current study results, accumulation of toxic metals was showed (33–34) vary according to participants' gender (36–37). An Italian study on children (aged 11–13 yr) reported higher levels of many elements (cadmium, cobalt, copper, zinc, and so on) in scalp hair of girls than in boys (38). In addition, many studies have revealed higher blood levels of manganese, copper, arsenic, cadmium, and selenium in females than males (39–41). Similar to hair and blood, a study has shown increased level of urinary cadmium in females than in males (35). On the other hand, some studies on children have reported no gender difference (42) or even lower blood levels of metal (lead) in girls than in boys (36,43,44). It is difficult to explain why metals levels were higher in girls than boys. However, it could be related to female vulnerability to absorb metals and/or a sex-related metabolic difference of metals (45), such as a higher bone release of metals (lead) in exchange with calcium, in female (46). In average, girls have a shorter growth period than boys due to complete their skeletal growth generally earlier and faster than boys (33, 39, 40).

In the present study, zinc showed a significant negative correlation with lead. Similar findings (in hair and blood) have been also reported in earlier studies (47–49). Perhaps, there is a competition between lead and zinc from absorption to the site of effects, as co-administration of zinc supple-

ments and lead in experimental study revealed a reduction in lead absorption and accumulation in organs (50–53). Therefore, zinc may have a protective effect against lead, mediating by competition to various binding sites (54), such as metal-binding proteins (e.g., metallothioneins) (55), and zinc allosteric sites (56).

The sources of metal exposure for the study participants were not known. However, hair concentrations of metals in the children may be affected by biological, personal, and environmental factors (57). Among these, environmental pollution (mainly anthropogenic) could be the main source of exposure in Tehran, as the city has considerable numbers of industrial sources. Although concentrations of many elements are lower in residential zones than in industrial areas, some elements are present in the same (arsenic and zinc) or even higher levels (manganese, antimony, and copper) in these zones (18). Similarly, children who live near a high traffic road were exposed to higher levels of metals than those who did not (58), but the present study information about participants' inhabitation did not include distance from roads. Some heavy metal contents were reported to be higher along the city roads (i.e., cadmium, 4 mg/kg; lead, 669 mg/kg; zinc, 614 mg/kg) (19) than the acceptable levels in natural soils (cadmium, 3 mg/kg; lead, 100 mg/kg; zinc, 300 mg/kg) (59). Therefore, the study participants may be exposed to potentially toxic metals mainly via industrial activities and vehicle exhaust fumes.

There were some limitations of the current study. First, the children's hair samples were collected at different ages (20–36 months old), while weight was taken from medical records from birth to 18 months old. Second, we did not wash the hair samples before analysis. Third, we had limited information on the children's diets and the fathers' anthropometric characteristics. The mothers were asked to provide details of the fathers' anthropometric characteristics, but the data were not sufficiently reliable to include in the statistical analyses. Finally, the current study did not measure metal levels in the tap water or in the air of Tehran.

Conclusion

Children with relatively low weight gain had higher level of cobalt, which the finding disappeared when adjusting for covariates. In addition, girls showed higher levels of toxic metals than in boys. This may suggest that gender plays an important role in the absorption and/or accumulation of metals. The study also revealed significant correlations among measured metals in the children scalp hair. Thus, the future studies should be conducted on mix metals set and should consider gender as an important factor in metals toxicity.

Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Acknowledgments

The authors would like to thank the research hospitals' staffs for gathering the participant's data, providing the medical records, contributing in sampling, and all lab measurements during the survey.

This study was supported in joint collaboration by 1) the Japanese National Institute for Occupational Safety and Health, 2) a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science, and 3) Tehran University of Medical Sciences.

Conflict of Interests

The authors declare that there is no conflict of interest.

References

1. Engle-Stone R, Ndjebayi AO, Nankap M, et al (2014). Stunting prevalence, plasma zinc concentrations, and dietary zinc intakes in a

- nationally representative sample suggest a high risk of zinc deficiency among women and young children in Cameroon. *J Nutr*, 144:382-91.
2. Jadhav SH, Sarkar SN, Patil RD, Tripathi HC (2007). Effects of subchronic exposure via drinking water to a mixture of eight water-contaminating metals: a biochemical and histopathological study in male rats. *Arch Environ Contam Toxicol*, 53:667-77.
3. Lech T (2002). Lead, copper, zinc, and magnesium content in hair of children and young people with some neurological diseases. *Biol Trace Elem Res*, 85:111-26.
4. McMichael AJ (1993). Lead and child development. *Arch Environ Health*, 48:125; author reply 126-7.
5. Fortoul TI, Moncada-Hernandez S, Saldivar-Osorio L, et al (2005). Sex differences in bronchiolar epithelium response after the inhalation of lead acetate (Pb). *Toxicology*, 207:323-30.
6. Jozefczak M, Remans T, Vangronsveld J, Cuypers A (2012). Glutathione Is a Key Player in Metal-Induced Oxidative Stress Defenses. *Int J Mol Sci*, 13:3145-75.
7. Sirivarasai J, Wananukul W, Kaojareen S, et al (2013). Association between Inflammatory Marker, Environmental Lead Exposure, and Glutathione S-Transferase Gene. *Biomed Res Int*, 2013:474963.
8. Boadi WY, Harris S, Anderson JB, Adunyah SE (2013). Lipid peroxides and glutathione status in human progenitor mononuclear (U937) cells following exposure to low doses of nickel and copper. *Drug Chem Toxicol*, 36:155-62.
9. Llorente-Cantarero FJ, Gil-Campos M, Benitez-Sillero Jde D, et al (2013). Profile of oxidant and antioxidant activity in prepubertal children related to age, gender, exercise, and fitness. *Appl Physiol Nutr Metab*, 38:421-6.
10. Hamon I, Valdes V, Franck P, et al (2011). [Gender-dependent differences in glutathione (GSH) metabolism in very preterm infants]. *Arch Pediatr*, 18:247-52.
11. Bellinger DC (1995). Interpreting the literature on lead and child development: the neglected role of the "experimental system". *Neurotoxicol Teratol*, 17:201-12.

12. Lavoie JC, Chessex P (1997). Gender and maturation affect glutathione status in human neonatal tissues. *Free Radic Biol Med*, 23:648-57.
13. Kelishadi R, Amiri M, Motlagh ME, et al (2014). Growth disorders among 6-year-old Iranian children. *Iran Red Crescent Med J*, 16:e6761.
14. Ahmadi A, Moazen M, Mosallaei Z, et al (2014). Nutrient intake and growth indices for children at kindergartens in Shiraz, Iran. *J Pak Med Assoc*, 64:316-21.
15. Kelishadi R, Ardalan G, Qorbani M, et al (2013). Methodology and Early Findings of the Fourth Survey of Childhood and Adolescence Surveillance and Prevention of Adult Non-Communicable Disease in Iran: The CASPIAN-IV Study. *Int J Prev Med*, 4:1451-60.
16. Payandeh A, Saki A, Safarian M, et al (2013). Prevalence of malnutrition among preschool children in northeast of Iran, a result of a population based study. *Glob J Health Sci*, 5:208-12.
17. Baghianimoghadam B, Karbasi SA, Golestan M, Kamran MH (2012). Determination of growth pattern of 7-12 years old children in YAZD city and comparison of it with WHO standards. *J Pak Med Assoc*, 62:1289-93.
18. Sekhvatjou MS, Hosseini Alhashemi A, Rostami A (2011). Comparison of trace element concentrations in ambient air of industrial and residential areas in Tehran city. *Biol Trace Elem Res*, 143:1413-23.
19. Saeedi M, Hosseinzadeh M, Jamshidi A, Pajoohehfar SP (2009). Assessment of heavy metals contamination and leaching characteristics in highway side soils, Iran. *Environ Monit Assess*, 151:231-41.
20. Soleimani S, Shahverdy MR, Mazhari N, et al (2014). Lead concentration in breast milk of lactating women who were living in Tehran, Iran. *Acta Med Iran*, 52:56-9.
21. Savabieasfahani M, Hoseiny M, Goodarzi S (2012). Toxic and essential trace metals in first baby haircuts and mother hair from Imam Hossein Hospital Tehran, Iran. *Bull Environ Contam Toxicol*, 88:140-4.
22. Serdar MA, Akin BS, Razi C, et al (2012). The correlation between smoking status of family members and concentrations of toxic trace elements in the hair of children. *Biol Trace Elem Res*, 148:11-7.
23. Mehra R, Juneja M (2003). Adverse health effects in workers exposed to trace/toxic metals at workplace. *Indian J Biochem Biophys*, 40:131-5.
24. Gil F, Hernandez AF, Marquez C, et al (2011). Biomonitorization of cadmium, chromium, manganese, nickel and lead in whole blood, urine, axillary hair and saliva in an occupationally exposed population. *Sci Total Environ*, 409:1172-80.
25. Klatka M, Blazewicz A, Partyka M, et al (2015). Concentration of Selected Metals in Whole Blood, Plasma, and Urine in Short Stature and Healthy Children. *Biol Trace Elem Res*, 166:142-8.
26. Skalnaya MG, Tinkov AA, Demidov VA, et al (2014). Hair toxic element content in adult men and women in relation to body mass index. *Biol Trace Elem Res*, 161:13-9.
27. Ahmed S, Rekha RS, Ahsan KB, et al (2013). Arsenic Exposure Affects Plasma Insulin-Like Growth Factor 1 (IGF-1) in Children in Rural Bangladesh. *PLoS One*, 8:e81530.
28. Saha KK, Engstrom A, Hamadani JD, et al (2012). Pre- and Postnatal Arsenic Exposure and Body Size to 2 Years of Age: A Cohort Study in Rural Bangladesh. *Environ Health Perspect*, 120:1208-14.
29. Wang SX, Wang ZH, Cheng XT, et al (2007). Arsenic and Fluoride Exposure in Drinking Water: Children's IQ and Growth in Shanyin County, Shanxi Province, China. *Environ Health Perspect*, 115:643-7.
30. Barbosa F, Jr., Ramires I, Rodrigues MH, et al (2006). Contrasting effects of age on the plasma/whole blood lead ratio in men and women with a history of lead exposure. *Environ Res*, 102:90-5.
31. Popovic M, McNeill FE, Chettle DR, et al (2005). Impact of occupational exposure on lead levels in women. *Environ Health Perspect*, 113:478-84.
32. Bjorkman L, Vahter M, Pedersen NL (2000). Both the environment and genes are important for concentrations of cadmium and lead in blood. *Environ Health Perspect*, 108:719-22.
33. Abdelouahab N, Mergler D, Takser L, et al (2008). Gender differences in the effects of organochlorines, mercury, and lead on

- thyroid hormone levels in lakeside communities of Quebec (Canada). *Environ Res*, 107:380-92.
34. Massanyi P, Tataruch F, Slameka J, et al (2003). Accumulation of lead, cadmium, and mercury in liver and kidney of the brown hare (*Lepus europaeus*) in relation to the season, age, and sex in the West Slovakian Lowland. *J Environ Sci Health A Tox Hazard Subst Environ Eng*, 38:1299-309.
 35. Jemal A, Graubard BI, Devesa SS, Flegal KM (2002). The association of blood lead level and cancer mortality among whites in the United States. *Environ Health Perspect*, 110:325-9.
 36. Duc Phuc H, Kido T, Dung Manh H, et al (2016). A 28-year observational study of urinary cadmium and β_2 -microglobulin concentrations in inhabitants in cadmium-polluted areas in Japan. *J Appl Toxicol*, 36:1622-1628.
 37. Chen J, Li M, Lv Q, et al (2015). Blood lead level and its relationship to essential elements in preschool children from Nanning, China. *J Trace Elem Med Biol*, 30:137-41.
 38. Varrica D, Tamburo E, Milia N, et al (2014). Metals and metalloids in hair samples of children living near the abandoned mine sites of Sulcis-Inglesiente (Sardinia, Italy). *Environ Res*, 134:366-74.
 39. Oulhote Y, Mergler D, Bouchard MF (2014). Sex- and age-differences in blood manganese levels in the U.S. general population: national health and nutrition examination survey 2011-2012. *Environ Health*, 13:87.
 40. Zhang LL, Lu L, Pan YJ, et al (2015). Baseline blood levels of manganese, lead, cadmium, copper, and zinc in residents of Beijing suburb. *Environ Res*, 140:10-7.
 41. Inoue Y, Umezaki M, Jiang H, et al (2014). Urinary concentrations of toxic and essential trace elements among rural residents in Hainan Island, China. *Int J Environ Res Public Health*, 11:13047-64.
 42. Allen Counter S, Buchanan LH, Ortega F (2015). Blood Lead Levels in Andean Infants and Young Children in Ecuador: An International Comparison. *J Toxicol Environ Health A*, 78:778-87.
 43. Xu J, Sheng L, Yan Z, Hong L (2014). Blood lead and cadmium levels in children: A study conducted in Changchun, Jilin Province, China. *Paediatr Child Health*, 19:73-6.
 44. Swaddiwudhipong W, Kavinum S, Papwijitsil R, et al (2014). Personal and environmental risk factors significantly associated with elevated blood lead levels in rural Thai children. *Southeast Asian J Trop Med Public Health*, 45:1492-502.
 45. Finley JW, Johnson PE, Johnson LK (1994). Sex affects manganese absorption and retention by humans from a diet adequate in manganese. *Am J Clin Nutr*, 60:949-55.
 46. Gulson B, Mizon K, Korsch M, Taylor A (2016). Revisiting mobilisation of skeletal lead during pregnancy based on monthly sampling and cord/maternal blood lead relationships confirm placental transfer of lead. *Arch Toxicol*, 90:805-16.
 47. Zaborowska W, Wiercinski J (1996). [Determination of lead, cadmium, copper and zinc in hair of children from Lublin as a test of environmental pollution]. *Rocz Panstw Zakl Hig*, 47:217-22.
 48. Noonan CW, Kathman SJ, Sarasua SM, White MC (2003). Influence of environmental zinc on the association between environmental and biological measures of lead in children. *J Expo Anal Environ Epidemiol*, 13:318-23.
 49. Kang-Sheng L, Xiao-Dong M, Juan S, et al (2015). Towards bio monitoring of toxic (lead) and essential elements in whole blood from 1- to 72-month old children: a cross-sectional study. *Afr Health Sci*, 15:634-40.
 50. Flora SJ, Kumar D, Das Gupta S (1991). Interaction of zinc, methionine or their combination with lead at gastrointestinal or post-absorptive level in rats. *Pharmacol Toxicol*, 68:3-7.
 51. Jamieson JA, Taylor CG, Weiler HA (2006). Marginal zinc deficiency exacerbates bone lead accumulation and high dietary zinc attenuates lead accumulation at the expense of bone density in growing rats. *Toxicol Sci*, 92:286-94.
 52. Batra N, Nehru B, Bansal MP (1998). The effect of zinc supplementation on the effects of lead on the rat testis. *Reprod Toxicol*, 12:535-40.
 53. Rowles TK, Womac C, Bratton GR, Tiffany-Castiglioni E (1989). Interaction of lead and zinc in cultured astroglia. *Metab Brain Dis*, 4:187-201.

54. Batra N, Nehru B, Bansal MP (2001). Influence of lead and zinc on rat male reproduction at 'biochemical and histopathological levels'. *J Appl Toxicol*, 21:507-12.
55. Goyer RA (1997). Toxic and essential metal interactions. *Annu Rev Nutr*, 17:37-50.
56. Guilarte TR, Miceli RC, Jett DA (1995). Biochemical evidence of an interaction of lead at the zinc allosteric sites of the NMDA receptor complex: effects of neuronal development. *Neurotoxicology*, 16:63-71.
57. Sukumar A (2002). Factors influencing levels of trace elements in human hair. *Rev Environ Contam Toxicol*, 175:47-78.
58. Rahbar MH, Samms-Vaughan M, Dickerson AS, et al (2015). Factors associated with blood lead concentrations of children in Jamaica. *J Environ Sci Health A Tox Hazard Subst Environ Eng*, 50:529-39.
59. Fabis W (1987) *Schadstoffbelastung Von Böden-Auswirkungen auf Böden-und Wassergalität Allg Festszeitsehr*. München: BLV Verlagsgesellschaft, pp. 128-131.