



## Evaluative Assay of Nuclear and Mitochondrial Genes to Diagnose *Leishmania* Species in Clinical Specimens

Ahmad Reza *ESMAEILI RASTAGHI*<sup>1</sup>, Adel *SPOTIN*<sup>1,2</sup>, Mohammad Reza *KHATAMI-NEZHAD*<sup>3</sup>, Mostafa *JAFARPOUR*<sup>3</sup>, Elnaz *ALAEENOVIN*<sup>1</sup>, Narmin *NAJAFZADEH*<sup>1</sup>, Neda *SAMEI*<sup>1,3</sup>, Neda *TALESHI*<sup>1,3</sup>, Somayeh *MOHAMMADI*<sup>1</sup>, \*Parviz *PARVIZI*<sup>1</sup>

1. Molecular Systematics Laboratory, Parasitology Department, Pasteur Institute of Iran, Tehran, Iran
2. Dept. of Medical Parasitology & Mycology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran
3. Dept. of Biology, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran

\*Corresponding Author: Email: [parp@pasteur.ac.ir](mailto:parp@pasteur.ac.ir)

(Received 18 Oct 2016; accepted 20 Jan 2017)

### Abstract

**Background:** Leishmaniasis as an emerging and reemerging disease is increasing worldwide with high prevalence and new incidence in recent years. For epidemiological investigation and accurate identification of *Leishmania* species, three nuclear and mitochondrial genes (ITS-rDNA, Hsp70, and Cyt *b*) were employed and analyzed from clinical samples in three important Zoonotic Cutaneous Leishmaniasis (ZCL) foci of Iran.

**Methods:** In this cross-sectional/descriptive study conducted in 2014-15, serous smears of lesions were directly prepared from suspected patients of ZCL in Turkmen in northeast, Abarkouh in center and Shush district in southwest of Iran. They were directly prepared from suspected patients and DNA was extracted. Two nuclear genes of ITS-rDNA, Hsp70 and one mitochondrial gene of Cyt *b* within *Leishmania* parasites were amplified. RFLP was performed on PCR-positive samples. PCR products were sequenced, aligned and edited with sequencer 4.1.4 and phylogenic analyses performed using MEGA 5.05 software.

**Results:** Overall, 203 out of 360 clinical samples from suspected patients were *Leishmania* positive using routine laboratory methods and 231 samples were positive by molecular techniques. *L. major*, *L. tropica*, and *L. turanica* were firmly identified by employing different molecular genes and phylogenic analyses.

**Conclusion:** By combining different molecular genes, *Leishmania* parasites were identified accurately. The sensitivity and specificity three genes were evaluated and had more advantages to compare routine laboratory methods. ITS-rDNA gene is more appropriate for firm identification of *Leishmania* species.

**Keywords:** *Leishmania* species, ITS-rDNA, Hsp70, Cyt *b*, Diagnosis, Iran

### Introduction

Leishmaniasis is one of the most important human protozoan parasitic diseases worldwide by increasing the prevalence and incidence rates in recent years (1-3). From three types of leishmaniasis in Iran, Zoonotic Cutaneous Leishmaniasis (ZCL) has a great distribution reported from more than half of Iranian provinces (4, 5). ZCL is a single cell parasitic disease that considered as a major health problem in many areas of Iran.

Rodents and other mammals are reservoir hosts and human infects the causative agent of *Leishmania major* accidentally by biting female sand flies (6, 7). The presence of ZCL in Turkmen Sahara (Golestan), Abarkouh (Yazd), and Shush (Khuzestan) has been demonstrated in addition new ZCL foci in some locations located in border of Iran and Iraq (7, 8). *L. major*, *L. turanica*, *L. jebilli* and a new *Leishmania* close to *L. jebilli* have been

reported in ZCL foci but *L. major* is the principal agent of ZCL in Iran (3, 9, 10). *Phlebotomus papatasi* is the main vector and the most important reservoir hosts are *Rhombomys opimus* and *Meriones libycus* in Turkmen Sahara and Abarkouh, and *Tatera indica* in Shush (8, 11-13).

ZCL represents a typical model of emerging and reemerging zoonosis disease (14). ZCL can cause substantial morbidity because of the presence of a chronic skin ulcer and the psychological effect of disfigurement (15). There are no proper vaccines to protect people against the parasites. Although, epidemiological investigation and finding *Leishmania* species are helpful in the diagnosis of ZCL; however they are not sufficient methods to identify firmly the agent of the disease. Accurate identification of the causative agent of *Leishmania* species is essential to give us knowledge of the *Leishmania* species in the endemic and specific geographical area, and better approach in control measurements and treatment of disease (16). Hence, firmly identification and determination of *Leishmania* parasites are indispensable advanced molecular methods for human cases because of being deficient information for routine laboratory techniques.

For this investigation, three nuclear and mitochondrial genes (ITS-rDNA, Hsp70, and Cyt *b*) of *Leishmania* were employed, compared and designed to identify the *Leishmania* species parasites circulating in suspected patients of ZCL in different regions in Northeast, Central and Southwest to Iran.

## Materials and Methods

In this cross-sectional/descriptive study conducted in 2014-2015, serous smears of lesions were directly prepared from suspected patients of ZCL in Turkmen Sahara (37° 13' 0" N 55° 0' 0" E) in northeast, Abarkouh (31° 7' 44.04" N 53° 16' 56.64" E) in center and Shush district (32° 11' 39.12" N 48° 14' 36.96" E) in southwest of Iran.

Human samples were collected from urban and rural areas surrounding Shush, Turkmen Sahara and Abarkouh districts of Iranian provinces. The personal information, lesion duration, type and number of lesion, ulcer(s)' location, patients' traveling to endemic regions and the grading numbers of amastigotes were recorded individually and kept confidential (Fig. 1).



Fig. 1: Iran map showing where sampling from suspected patients of ZCL took place


Giemsa Stained slides were prepared. Amastigote presence in slides was observed and graded under light microscope (Table 1). DNA was extracted from the slides using Phenol-Chlorophorm method with minor modification (3).

Two nuclear genes of ITS-rDNA (480 bp), Hsp70 (750 bp) and one mitochondrial gene of Cyt *b* (880 bp) within *Leishmania* parasites were amplified to detect any *Leishmania* infection among samples from suspected patients following the primers and protocols (16).

To select suitable restriction enzyme, CLC DNA workbench 5.2 software (CLC bio A/S, Aarhus, Denmark) In Silico was employed. By choosing sequences of different *Leishmania* species regis-

tered in GenBank; enzyme *BsuRI* (HaeIII) for ITS-rDNA and *SspI* enzyme for Cyt *b* gene were selected (Fig. 2). Hsp70 gene was not used for RFLP. Selected enzyme had different cut sites at GG↓CC in various species of *Leishmania* as a proper enzyme for PCR product digestion. RFLP was performed on PCR-positive samples for identification of *Leishmania* species. A master mix containing enzyme, Buffer and PCR product was prepared and stored at 37 °C for 4 hours.

Moreover, for accurate identification and confirmation of the specific PCR products were sequenced, aligned and edited with sequencer 4.1.4 software and phylogenetic analyses was done using MEGA 5.05 software (Fig. 3).




**Fig. 2:** CLC DNA workbench software showing different digestion sites of *BsuRI* and *SspI* enzymes affecting ITS-rDNA, Cyt *b*, and HSP70 genes In Sillico

## Results

The serous smears of lesions from 360-suspected patients (clinical samples) were sampled. Using routine laboratory methods, 203 samples from suspected patients were identified positive (Table 1). By employing molecular tools, using three different genes, 231 samples were *Leishmania* posi-

tive (Table 2, Fig. 2). After RFLP and/or sequencing; *L. major*, *L. tropica* and *L. turanica* were firmly identified (Table 2).

Shush district in south of Iran had higher infections than two other location where were sampled (87/360). The highest infection was in age group 10-25 yr (40/360) and male had more positive than females (Table 1).



**Fig. 3:** Maximum Likelihood algorithm showing the haplotypes of Cyt b gene for the isolates of Leishmania species using MEGA.5.05

Seventy-nine percent of the lesions were wet, 40% of them were in the hands, 42% of patients had single lesions and 32% of the slides were 4+, which was the highest velocity among the surveyed categories (Table 1).

Overall, 231 (64.1%) of samples were confirmed to be *Leishmania* positive via molecular methods (Table 2), of which 46.7%, 23.8%, and 28.5% were tested positive with by ITS-rDNA, Hsp70 and Cyt b genes, respectively (Table 2).

After RFLP and/or sequencing; from 231 *Leishmania* positive samples 217 (93.9%), 4 (1.73%) and 2 (0.86%) were indefinitely identified as *L. major*, *L. tropica*, and *L. turanica*, respectively. In addition, 8 (3.46%) of the samples were unable to

be accurately confirmed due to lack of PCR product and/or bad sequence reads (Table 2).

In this investigation, *L. major*, *L. tropica* and *L. turanica* were unambiguously typed and identified after analyzing and sequencing with molecular software with comparison to those sequences which have already been registered in GenBank in case of any similarity and homology. The obtained sequences of this investigation were homolog with *Leishmania* species after direct sequencing, editing, aligning and comparing with the sequences submitted to GenBank using Sequencher™ 4.1.4.

In the present work, we have compared the specificity and sensitivity of three nuclear and mitochondrial genes, we discovered that sensitivity of ITS-rDNA was more than Cyt b and Hsp70 and ITS-rDNA show the highest specificity for *Leishmania* species.

## Discussion

In this investigation, *L. major*, *L. tropica* and *L. turanica* among suspected patients were firmly identified in three endemic ZCL foci, Iran. Three well-known molecular markers ITS-rDNA, Hsp70 and Cyt b genes were employed (3). Three genes were applied concurrently to compare specificity, sensitivity and to increase the chance of detecting any *Leishmania* parasites even in low concentrations (17).

ITS-rDNA gene is a nuclear, linear, homogenous and conserves gene with low intracellular polymorphism and readable sequences. ITS-rDNA is ideal for phylogenetic analysis (12).

Cyt b as a mitochondrial marker can identify the novel nucleotide variations (haplotype) superior to ITS-rDNA (nucleus gene), this is associated with highly being conserved and high copy numbers of Cyt b per cell (20–50 maxi circles in 30 Kbp) (Fig. 2) (5, 18, 19).

ITS-rDNA shows high sensitivity because of approximately 20–400 copies of gene for *Leishmania* but not for differentiate *Leishmania* species. Cyt b as an evolutionary mitogenome marker has its semi conserved structure and low copy number

could be able to utilize in the discrimination of new mutants, whereas no significant mutant was observed in ITS-rDNA sequences (7).

By analyzing of CLC DNA workbench software, digestion sites of different enzymes were recognized on ITS-rDNA, Hsp70 and Cyt b genes of *Leishmania* species (Fig. 2). *BsuRI* enzyme has one digestion site in *L. major* for ITS-rDNA gene that gives two fragments (120bp and 300 bp), three digestion sites for Hsp70 gene. *SspI* enzyme has one digestion site (~ 500bp) for Cyt b gene in *L. major*. The different and variation fragments by digesting site make RFLP method, which generated species-specific patterns of bands visualized in agarose gels, a useful technique for accurate determination of *Leishmania* species.

Only amplifying DNA by PCR and observation of relevant band in agarose gels without sequences, molecular and phylogenetic analysis could not be effective and trusted firmly identification of *Leishmania* species (20).

Regarding data of this investigation, males are at higher risk than females for *Leishmania* infection in Iran because of wearing Hejab, covering skin by females, preventing sand fly bites and decreasing the risk for leishmaniasis. Moreover, men usually work in farms, fields and mostly sleep outside during the adult sand flies activity; these provide a good source of blood meals and transferring *Leishmania* parasites in ZCL foci (21). These also could be reason for high rate of infection in 10-25 age groups.

## Conclusion

Different genes combined for accurate identifications of clinical samples of *Leishmania* parasites sampled in three well-known ZCL foci in Iran. To find standardized, sensitive, specific, practical and reproducible genes for molecular identification and typing *Leishmania* species were evaluated and compared using three genes in this investigation. Molecular tools are more trustable than routine laboratory methods, and ITS-rDNA gene is more appropriate for accurate identification of *Leishmania* species.

## Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

## Acknowledgments

Pasteur Institute of Iran supported the work; grant 501 awarded to Dr. Parviz Parvizi. The authors would like to thank Health Care authorities and patients for their help and support during this study. We thank Mehdi Baghban for helping with the field working in Molecular Systematic Laboratory. A part of this research through MSc studentships to Neda Samei based at the Pasteur Institute of Iran, and registered for Islamic Azad University, Tonekabon, Iran.

## Conflict of Interest

The authors declare that there is no conflict of interest.

## References

1. Zangger H, Ronet C, Desponds C et al (2013). Detection of *Leishmania* RNA virus in *Leishmania* parasites. *PLoS Negl Trop Dis*, 7(1):e2006.
2. Vilas VJ, Maia-Elkhoury AN, Yadon ZE, Cosivi O, Sanchez-Vazquez MJ (2014). Visceral leishmaniasis: a One Health approach. *Vet Rec*, 175(2):42-4.
3. Bordbar A, Parvizi P (2014). High infection frequency, low diversity of *Leishmania major* and first detection of *Leishmania turanica* in human in northern Iran. *Acta Trop*, 133:69-72.
4. Yaghoobi-Ershadi MR (2012). Phlebotomine sand flies (Diptera: Psychodidae) in Iran and their role on *Leishmania* transmission. *J Arthropod Borne Dis*, 6(1):1-17.
5. Spotin A, Rouhani S, Parvizi P (2014). The associations of *Leishmania major* and *Leishmania tropica* aspects by focusing their morphological and molecular features on clinical appearances

- in Khuzestan province, Iran. *Biomed Res Int*, 913510. doi: 10.1155/2014/913510.
6. Parvizi P, Ready PD (2008). Nested PCRs and sequencing of nuclear ITS-rDNA fragments detect three *Leishmania* species of gerbils in sandflies from Iranian foci of zoonotic cutaneous leishmaniasis. *Trop Med Int Health*, 13(9):1159-71.
  7. Spotin A, Rouhani S, Ghaemmaghami P A, et al (2015). Different Morphologies of *Leishmania major* Amastigotes with No Molecular Diversity in a Neglected Endemic Area of Zoonotic Cutaneous Leishmaniasis in Iran. *Iran Biomed J*, 19(3):149-59.
  8. Najafzadeh N, Sedaghat MM, Sultan SS et al (2014). The existence of only one haplotype of *Leishmania major* in the main and potential reservoir hosts of zoonotic cutaneous leishmaniasis using different molecular markers in a focal area in Iran. *Rev Soc Bras Med Trop*, 47(5):599-606.
  9. Parvizi P, Mauricio I, Aransay AM, Miles MA, Ready PD (2005). First detection of *Leishmania major* in peridomestic Phlebotomus papatasi from Isfahan province, Iran: comparison of nested PCR of nuclear ITS ribosomal DNA and semi-nested PCR of minicircle kinetoplast DNA. *Acta Trop*, 93(1):75-83.
  10. Zahraei-Ramazani A, Kumar D, Mirhendi H et al (2015). Morphological and genotypic variations among the species of the subgenus Adlerius (Diptera: Psychodidae, Phlebotomus) in Iran. *J Arthropod Borne Dis*, 9(1):84-97.
  11. Jafari R, Najafzadeh N, Sedaghat MM, Parvizi P (2013). Molecular characterization of sandflies and *Leishmania* detection in main vector of zoonotic cutaneous leishmaniasis in Abarkouh district of Yazd province, Iran. *Asian Pac J Trop Med*, 6(10):792-7.
  12. Mirzaei A, Rouhani S, Taherkhani H et al (2011). Isolation and detection of *Leishmania* species among naturally infected *Rhombomys opimus*, a reservoir host of zoonotic cutaneous leishmaniasis in Turkmen Sahara, North East of Iran. *Exp Parasitol*, 129(4):375-80.
  13. Mohammadi S, Parvizi P (2016). Simultaneous Morphological and Molecular Characterization of *Tatera indica* in Southwestern Iran. *J Arthropod Borne Dis*, 10(1):55-64.
  14. Ashford RW (2000). The leishmaniasis as emerging and reemerging zoonoses. *Int J Parasitol* 30(12):1269-81.
  15. Bern C, Maguire JH, Alvar J (2008). Complexities of assessing the disease burden attributable to leishmaniasis. *PLoS Negl Trop Dis*, 2(10):e313.
  16. Tellevik MG, Muller KE, Løkken KR, Nerland AH (2014). Detection of a broad range of *Leishmania* species and determination of parasite load of infected mouse by real-time PCR targeting the arginine permease gene AAP3. *Acta tropica*, 137:99-104.
  17. Bensoussan E, Nasereddin A, Jonas F, Schnur LF, Jaffe CL (2006). Comparison of PCR assays for diagnosis of cutaneous leishmaniasis. *Int J Parasitol* 44(4):1435-9.
  18. Degli Esposti M, De Vries S, Crimi M, Ghelli A, Patarnello T, Meyer A (1993). Mitochondrial cytochrome b: evolution and structure of the protein. *Biochim Biophys Acta*, 1143(3):243-71.
  19. Howell N, Gilbert K (1988). Mutational analysis of the mouse mitochondrial cytochrome b gene. *J Mol Biol*, 203(3):607-17.
  20. Schonian G, Nasereddin A, Dinse N et al (2003). PCR diagnosis and characterization of *Leishmania* in local and imported clinical samples. *Diagn Microbiol Infect Dis*, 47(1):349-58.
  21. Leishmaniasis WHOECotCot, Meeting WHOECotCotL, World Health O. Control of the Leishmaniasis: Report of a WHO Expert Committee: World Health Organization; 1990.

**Table 1:** *Leishmania* identifications from suspected patients of three endemic ZCL foci, Iran

| Sampling site |                | Sex       |                     | Grading positive slides |                         |      |        |                                   |                                  | Lesion form                     |                             | Lesion No.                    |                             |     | Lesion Site |        |        |      |      |      |              |       |  |  |
|---------------|----------------|-----------|---------------------|-------------------------|-------------------------|------|--------|-----------------------------------|----------------------------------|---------------------------------|-----------------------------|-------------------------------|-----------------------------|-----|-------------|--------|--------|------|------|------|--------------|-------|--|--|
| Provinces     | Districts      | Age Group | Positive sample No. | Suspected patients NO   | Microscopic Observation | Male | Female | +1<br>1-10 Parasite / 1000 fields | +2<br>1-10 Parasite / 100 fields | +3<br>1-10 Parasite / 10 fields | +4<br>1-10 Parasite / field | +5<br>10-100 Parasite / field | +6<br>>100 Parasite / field | Wet | Dry         | Single | Double | Many | Hand | Foot | Face & Noose | Other |  |  |
| Yazd          | Abarkouh       | 1-5       | 5                   | 90                      | 44                      | 31   | 20     | 5                                 | 6                                | 9                               | 14                          | 13                            | 3                           | 41  | 10          | 30     | 15     | 16   | 26   | 12   | 7            | 6     |  |  |
|               |                | 5-10      | 15                  |                         |                         |      |        |                                   |                                  |                                 |                             |                               |                             |     |             |        |        |      |      |      |              |       |  |  |
|               |                | 10-25     | 20                  |                         |                         |      |        |                                   |                                  |                                 |                             |                               |                             |     |             |        |        |      |      |      |              |       |  |  |
|               |                | >25       | 10                  |                         |                         |      |        |                                   |                                  |                                 |                             |                               |                             |     |             |        |        |      |      |      |              |       |  |  |
| Golestan      | Turkmen Sahara | 1-5       | 10                  | 150                     | 80                      | 61   | 29     | 6                                 | 10                               | 12                              | 28                          | 24                            | 10                          | 73  | 17          | 32     | 34     | 24   | 36   | 29   | 16           | 9     |  |  |
|               |                | 5-10      | 25                  |                         |                         |      |        |                                   |                                  |                                 |                             |                               |                             |     |             |        |        |      |      |      |              |       |  |  |
|               |                | 10-25     | 30                  |                         |                         |      |        |                                   |                                  |                                 |                             |                               |                             |     |             |        |        |      |      |      |              |       |  |  |
|               |                | >25       | 25                  |                         |                         |      |        |                                   |                                  |                                 |                             |                               |                             |     |             |        |        |      |      |      |              |       |  |  |
| Khuzestan     | Shush          | 1-5       | 8                   | 120                     | 79                      | 46   | 44     | 5                                 | 8                                | 14                              | 30                          | 25                            | 8                           | 69  | 21          | 35     | 25     | 20   | 31   | 39   | 10           | 10    |  |  |
|               |                | 5-10      | 18                  |                         |                         |      |        |                                   |                                  |                                 |                             |                               |                             |     |             |        |        |      |      |      |              |       |  |  |
|               |                | 10-25     | 40                  |                         |                         |      |        |                                   |                                  |                                 |                             |                               |                             |     |             |        |        |      |      |      |              |       |  |  |
|               |                | >25       | 24                  |                         |                         |      |        |                                   |                                  |                                 |                             |                               |                             |     |             |        |        |      |      |      |              |       |  |  |
| Total         |                |           | 231/360<br>(64.1%)  | 203                     |                         | 231  |        | 16                                | 24                               | 35                              | 72                          | 62                            | 21                          | 183 | 48          | 97     | 74     | 60   | 93   | 80   | 33           | 25    |  |  |
|               |                |           |                     |                         |                         |      |        | 7%                                | 10%                              | 15%                             | 32%                         | 27%                           | 9%                          | 79% | 21%         | 42%    | 32%    | 26%  | 40%  | 35%  | 14%          | 11%   |  |  |
|               |                |           |                     |                         |                         |      |        | 231                               |                                  |                                 |                             |                               | 231                         |     |             |        |        | 231  |      |      |              |       |  |  |

**Table 2:** Different *Leishmania* species infection detected from suspected patients of ZCL from three endemic foci of Iran based on molecular markers

| Provinces | Collection site |                            | Microscopic Observation | Sex     |        | Positive samples using molecular tools | Leishmania positive via amplified genes |               |               | Molecular Methods                   |                   |                    |                |
|-----------|-----------------|----------------------------|-------------------------|---------|--------|--|---|---------------|---------------|-------------------------------------|-------------------|--------------------|----------------|
|           | Districts       | Villages                   |                         | Male    | Female |  | ITS-rDNA                                | Hsp 70        | Cyt b         | RFLP with <i>BsuRI</i> & sequencing |                   |                    |                |
|           |                 |                            |                         |         |        |  |   |               |               | <i>L. major</i>                     | <i>L. tropica</i> | <i>L. turanica</i> | Non-identified |
| Yazd      | Abarkouh        | Abarghasr                  | 44                      | 20      | 31     | 51/90                                  | 30                                      | 10            | 11            | 48                                  | 4                 | 0                  | 2              |
|           |                 | Chahgir                    |                         |         |        |  |   |               |               |                                     |                   |                    |                |
|           |                 | Harooni Abarkouh           |                         |         |        |  |   |               |               |                                     |                   |                    |                |
| Golestan  | Turkmen Sahara  | Kooran                     | 80                      | 29      | 61     | 90/150                                 | 45                                      | 20            | 20            | 85                                  | 0                 | 2                  | 3              |
|           |                 | Dashboroun Gharegol Hootan |                         |         |        |  |   |               |               |                                     |                   |                    |                |
| Khuzestan | Shush           | Haft Tape                  | 79                      | 44      | 46     | 90/120                                 | 35                                      | 25            | 25            | 87                                  | 0                 | 0                  | 3              |
|           |                 | Sorkhe Aljazayer Banader   |                         |         |        |  |   |               |               |                                     |                   |                    |                |
|           | Total           |                            | 203/360                 | 93      | 138    | 231/360                                | 110/231 (47.6)                          | 55/231 (23.8) | 66/231 (28.5) | 217/231 (93.9%)                     | 4/231 (1.73%)     | 2/231 (0.86%)      | 8/231 (3.46%)  |
|           |                 |                            |                         | 231/360 |        |  | 231/360                                 |               |               |                                     |                   |                    |                |