



Increases of Obesity among Iranian Adults: An Alarm for Policymakers

*Salman KHAZAEI¹, Saeid BASHIRIAN², *Ensiyeh JENABI³*

1. Research Center for Health Sciences, Hamadan University of Medical Sciences, Hamadan, Iran
2. Social Determinants of Health Research Center, Hamadan University of Medical Sciences, Hamadan, Iran
3. Autism Spectrum Disorders Research Center, Hamadan University of Medical Sciences, Hamadan, Iran

*Corresponding Author: Email: en.jenabi@yahoo.com

(Received 12 Jan 2019; accepted 24 Jan 2019)

Dear Editor in Chief

Obesity is one of the major public health problems of the twenty-first century (1). The continuation of obesity lead to increased morbidity risk in later life such as diabetes, coronary heart disease (CHD) and some of cancers (1-3). The prevalence of obesity has been elevating rapidly all round the world and is estimated to elevate further (4). Weight increases from imbalance of positive energy that calories expended more than those consumed to sup-

port basal metabolism, activity and growth, although the etiology of obesity is multi-factorial (5). Iran is one of the countries with the high prevalence of obesity among adults. Therefore, we elevated the prevalence and trend of obesity among Iranian adults (6).

Figure 1 shows the prevalence of obesity among adults in Iran compared the other WHO regions in 2016.




Fig. 1: Prevalence of obesity among adults (%), BMI \geq 30, age-standardized estimates by WHO region in 2016

The highest prevalence of obesity among both males and females are found in Americas region (25.9% and 31% respectively). While South East Asia countries with the 6.1% prevalence rate for females and 3.3% for males had the lowest rate. The prevalence rate of obesity in Iran was higher than all regions for females (32.2%). Except Americas region and prevalence rate of obesity for Iranian men was higher than other WHO regions (19.3%), which was much higher than global average (15.1% for females and 11.1% for males) for both males and females.

As shown in Fig. 2, trend of obesity for both males and females was increasing from 2000 to 2016. For

males, prevalence of obesity among adults reached from 10.3% in 2000 to 19.3% in 2016 (Annual percent change (APC): 3.94%, $P < 0.05$), while for females reached from 19.3% in 2000 to 32.2% in 2016 (Annual percent change (APC): 2.14%, $P < 0.05$). The reason of obesity among Iranian adults is patterns of inappropriate dietary habits, using of junk food and limitation of physical activity due to society's modernization. There is a need to establish programs for elevating awareness among the population in related to the obesity complications, control methods of obesity and performing of organized educational and preventive programs in adults.




Fig. 2: Trend analysis for prevalence of obesity among Iranian adults (BMI \geq 30) (2000-2016)

Conflict of interest

None declare.

References

1. Simmonds M, Burch J, Llewellyn A et al (2015). The use of measures of obesity in childhood for predicting obesity and the development of obesity-related diseases in adulthood: a systematic review and meta-analysis. *Health Technol Assess*, 19(43):1-336.
2. Jenabi E, Poorolajal J (2015). The effect of body mass index on endometrial cancer: a meta-analysis. *Public Health*, 129(7):872-80.
3. Poorolajal J, Jenabi E (2016). The association between BMI and cervical cancer risk: a meta-analysis. *Eur J Cancer Prev*, 25(3):232-8.
4. Low S, Chin MC, Deurenberg-Yap M (2009). Review on epidemic of obesity. *Ann Acad Med Singapore*, 38(1):57-9.
5. Suarez MA, Crinion KM (2015). Food choices of children with autism spectrum disorders. *International Journal of School Health*, 2(3): e27502.
6. World Health Organization. Prevalence of obesity among adults: Country Profiles 2016. 2016 [cited 10 Aug 2018]; http://www.who.int/gho/ncd/risk_factors/overweight_obesity/obesity_adults/en/