



Phylogeography and Genetic Diversity of Human Hydatidosis in Bordering the Caspian Sea, Northern Iran by Focusing on *Echinococcus granulosus* Sensu Stricto Complex

Abolghasem SIYADATPANAH^{1,2}, Ahmad DARYANI³, Shahabeddin SARVI³, Adel SPOTIN⁴, Mehdi SHARIF³, Reza ESMAEELZADEH DIZAJI⁵, Davood ANVARI³, Amir EMAMI ZEYDI⁶, Mohammad Hasan KOHANSAL⁷, Nelson Ivan AGUDELO HIGUITA⁸, Seyed Abdollah HOSSEINI³, Mojgan ARYAEIPOUR⁹, *Shirzad GHOLAMI³

1. Ferdows School of Paramedical and Health, Birjand University of Medical Sciences, Birjand, Iran
2. Student Research Committee, Mazandaran University of Medical Sciences, Sari, Iran
3. Department of Parasitology, Toxoplasmosis Research Center, Mazandaran University of Medical Sciences, Sari, Iran
4. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran
5. Department of Poultry Disease, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran
6. Department of Medical-Surgical Nursing, Nasibeh School of Nursing and Midwifery, Mazandaran University of Medical Sciences, Sari, Iran
7. School of Medicine, Bam University of Medical Sciences, Bam, Iran
8. Division of Infectious Diseases Department of Medicine, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma, USA
9. Department of Medical Parasitology and Mycology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

*Corresponding Author: Email: sgholami200@gmail.com

(Received 09 Mar 2019; accepted 20 May 2019)

Abstract

Background: Human *Echinococcosis* is a cyclo-zoonotic infection caused by tapeworms of the *Echinococcus granulosus* sensu stricto complex. The detection of mitochondrial genome data of genus *Echinococcus* can reflect the taxonomic status, genetic diversity, and population structure genetics.

Methods: Totally, 52 formalin-fixed paraffin-embedded (FFPE) tissue samples from patients with histologically confirmed CE were collected from Mazandaran province, Iran in the period of Mar 1995 to May 2018. All extracted DNAs from (FFPE) tissue samples were subjected to amplify by polymerase chain reactions method targeting cytochrome c oxidase subunit 1 (*cox1*) gene. All PCR amplicons were sequenced to phylogenetic analysis and genetic diversity.

Results: Molecular analysis showed that 50(96.1%) and 2 (3.84%) isolates were identified as G1 and G3 *E. granulosus* genotypes, respectively. DNA sequence analyses indicated a high gene diversity for G1 (Haplotype diversity: 0.830) and G3 genotypes (Hd: 1.00). Based on multiple sequence alignment analyses, 7 (13.46%; G1 genotype) and 2 (3.84%; G3 genotype) new haplotypes were unequivocally identified.

Conclusion: G3 genotype (Buffalo strain) was identified from two human hydatidosis isolates in the region. Present study strengthens our knowledge about taxonomic status, transmission patterns of *Echinococcus* parasite to human and heterogeneity aspects of this parasite in clinical CE isolates of Northern Iran.

Keywords: Human hydatidosis; Genetic diversity; Phylogeography; Iran



Introduction

Cystic *Echinococcosis* (CE) caused by metacestode *E. granulosus* genotypes remains a potential neglected problem global for matters relating to zoonotic disease and the economic loss (1,2). About 58% of the total populations of Central Asian countries including Iran, Kazakhstan, Tajikistan, Turkmenistan, Uzbekistan, Afghanistan, Mongolia, Pakistan, and Western China are potentially at risk for this orphan disease (3).

Many factors contribute to the transmission and development of CE in humans and other animals, the most important of which are age, occupation, location (urban or village), parasite genotype, and definitive host (dogs or other carnivores) populations in the region (4-7). Human beings are the incidental intermediate host of the tapeworm *E. granulosus*, the larvae of which are forming anywhere from the human body (8-10).

Based on phylomolecular analyses of *Echinococcus* mitochondrial markers (e.g., *cox1/nad1*), ten strains which include *E. granulosus sensu stricto* (s.s.) (genotypes G1–G3), *E. equinus* (genotype G4), *E. ortleppi* (genotype G5), and *E. canadensis* (genotypes G6/G7, G8, G10) (11,12). *E. granulosus* (G1) is predominantly addressed as widespread distribution in Iran, where the genotypes G1, G2, G3, G5, G6, and G7 have sympatrically been identified from hyperendemic foci of Iran (13-15). Although, there is no community based large screening studies in Iran, based on hospital records the prevalence rate has been estimated to be 1.18-3 per 100,000 populations (16). One of the neglected hyperendemic regions of CE with various ecosystem zones is situated in bordering the Caspian Sea, Mazandaran Province, Northern Iran where there is no comprehensive phylomolecular data on structure of *Echinococcus* parasite in human hydatidosis. About 79 patients of CE surgeries were performed for ten years from 2005 to 2015 in this region. Identification of genetic features and emergent novel haplotypes of *E. granulosus* isolates in human can explicitly reflect new insights of cross-transmission patterns of *Echinococ-*

cus among canids-intermediate hosts, epizootiology, gene migration of parasite, evolutionary patterns, and circulation of probable drug-resistance alleles (17,18).

We aimed to assess phylogeography and inter-intra genetic variation levels of *Echinococcus* spp., from clinical samples of hospitalized patients in bordering the Caspian Sea, Northern Iran based on sequencing of cytochrome c oxidase I (*cox1*).

Materials and Methods

Study area and Sample collection

Mazandaran Province is located on a South-East coast of the Caspian Sea between 36.2262° N, 52.5319° E and covers an area of 23,842 km². The area has a temperate climate with 70%-100% relative humidity. The average annual temperature and rainfall ranges from 10–35 °C and 800-1,200 mm, respectively. It has various ecosystems including grasslands, sea and forests the province consists of 15 counties and has a population of around 2,922,432 inhabitants which living in rural and urban areas (19). Fifty-two (n=52 including 30 females) tissue samples (FFPE; Formalin-Fixed, Paraffin-Embedded) of humans were collected from the archives of different hospitals in Mazandaran Province including Amol, Babol, Chalus, Qaemshahr, Neka, and Sari hospitals in which CE patients had undergone surgery between Mar 1995 to May 2018 (Fig. 1).

DNA extraction and PCR procedure

To extract the genomic DNA of human tissue samples, 10 to 12 μm thick sections were obtained from FFPE blocks with microtome blades, were transferred to a sterile tube separately for each sample, for deparaffinized, 1 ml xylene was added to sections for 12 min at 37 °C then centrifuged at 1500 g for 5 min, and the supernatant was discarded. According to the protocol (20) for rehydration, sections were followed in 100%, 90%, 80%, and 70% ethanol, after that, the 70% ethanol was omitted and added lysis buffer.




Fig. 1: Map of Iran presenting study counties in bordering the Caspian Sea, Northern Iran

The specimens were digested 24 h, at 56 °C within 400 µL of the lysis buffer with the addition of 40 µL proteinase K (20 mg/ml). Genomic DNA extraction of each sample was done by (WizPrep™ gDNA Mini Kit (Cell/Tissue) according to the manufacturing company (wizbiosolution/ Korea) then the genomic DNA was kept at -20 °C until further use in PCR. The *cox1* gene was amplified with forward and reverse primers JB3-5'-TTTTTTT-GGGCATCCTGAGGTTTAT-3', JB4.5-5'-TAAAGAAAGAACATAATGAAAATG-3', respectively. The PCR reaction was used to amplify the *cox1* gene by 35 cycles 94 °C for 5 min as an initial denaturation, 94 °C for 30 sec, 50 °C for 45 sec, 72 °C for 35 sand a final extension at 72 °C for 10 min. The amplification products PCR were run on a 1.5% agarose gel electrophoresis (21).

Mitochondrial DNA sequencing, haplotype network, and phylogenetic tree

All PCR amplicons were sequenced by targeting *cox1* gene. To show the genealogical relationships of *Echinococcus* mutations, a haplotype network was constructed based on Median Joining algorithm (PopART software) (22). According to the analysis of molecular variance (AMOVA), the diversity indices (Nucleotide diversity (Nd; π) and Haplotype

(gene) diversity (Hd) and neutrality indices (Fu's Fs statistic and Tajima's D) were assessed by using DnaSP software version 5.10 (23). The divergence and percent identity among the edited sequences of *Echinococcus* genotypes were drawn using the MegAlign program. To reveal taxonomic status of *Echinococcus* strains, a phylogenetic tree was made based on maximum likelihood algorithm with Kimura 2-parameter (MEGA 5.05 software). Multiple alignments among the DNA sequences and amino-acid sequences were performed based on Clustal W method (BioEdit software, ver. 7.0.5). This research has been approved by the Deputy of research and technology, Mazandaran University of Medical Sciences Ethics Committee with ethical standards a special code: IR.MA-ZUMS.REC.96.2929.

Results

Demographic data, PCR, sequence analyses, and pairwise distance matrix

Overall, 52 human patients infected with hydatid cysts were collected from 11 counties of Mazandaran Province. Demographic data, identified strain, collected county names, and location of cystectomy organs are fully given in Table 1.

Table 1: Demographic data, identified strain, collected county names and location of cyst (Organs) in *Cystic Echinococcus* patients (n= 52) from Apr 2017 to May 2018 in Mazandaran province, Northern Iran

<i>Variable</i>	<i>Number of patients (%)</i>	<i>Genotype G1 (%)</i>	<i>Genotype G3 (%)</i>
<i>Age group(years)</i>			
10–20	1 (1.92)	1 (1.92)	-
>20–30	5 (9.62)	5 (9.62)	-
>30–40	12 (23.08)	12 (23.08)	-
>40–50	10 (19.23)	10 (19.23)	-
>50–70	24 (46.15)	22 (42.30)	2 (3.85)
Sex			
Male	22 (42.30)	22 (42.30)	-
Female	30 (57.70)	28 (53.84)	2 (3.85)
Cyst Location			
Liver	31 (59.62)	29 (55.76)	2 (3.85)
Lung	16 (30.77)	16 (30.77)	-
Kidney	2 (3.85)/	2 (3.85)	-
Ovary	1 (1.92)	1 (1.92)	-
Bile duct	1 (1.92)	1 (1.92)	-
Spleen	1 (1.92)	(1.92)	-
Residence			
City	18 (34.62)	17 (32.69)	1 (1.92)
Village	34 (65.38)	33 (63.46)	1 (1.92)
County names			
Amol	13 (25.00)	13 (25.00)	-
Babol	14 (26.92)	14 (26.92)	-
Babolsar	1 (1.92)	1 (1.92)	-
Behshahr	1 (1.92)	1 (1.92)	-
Tonekabon	2 (3.85)	2 (3.85)	-
Chalus	3 (5.77)	2 (3.85)	1 (1.92)
Ramsar	2 (3.85)	2 (3.85)	-
Sari	1 (1.92)	1 (1.92)	-
FereidunKenar	1 (1.92)	1 (1.92)	-
QaemShahr	1 (1.92)	1 (1.92)	-
MahmudAbad	2 (3.85)	2 (3.85)	-
Neka	1 (1.92)	1 (1.92)	-
Nur	6 (11.54)	6 (11.54)	-
Noshahr	4 (7.69)	3 (5.76)	1 (1.92)

The age ranges of the patients were between 20–68 (average 42) yr old. The amplified PCR products of the *Cox1* gene (444bp) were sequenced for 52 isolates. Overall, 50 (96.1%) and 2 (3.84%) isolates were found to correspond to *E. granulosus* G1 and G3 genotypes, respectively. All of the G3 genotype (n=2; 3.85%) were only found at liver hydatid cyst, while the G1 genotype was characterized

by hydatid cysts of liver, lungs, kidney, ovary, bile duct, and spleen (Table 1). Based on multiple sequence alignment analyses, 7 (13.46%; G1 genotype) and 2 (3.84%; G3 genotype) new haplotypes were unequivocally identified (Table 2). Distribution of identified haplotypes of *E. granulosus* G1 and G3 genotypes based on *cox1* registered sequences is given in Table 3.

Table 2: Diversity indices of *E.granulosus* G1 and G3 genotypes based on cytochrome c oxidase subunit 1 mitochondrial sequence in Mazandaran Province, Northern Iran

Parasite	Diversity indices				Neutrality indices		
	N	Nh (%)	Hd± SD	Nd (π)	No. of polymorphic sites	Tajima's D*	Fu's Fs statistic**
<i>E. granulosus</i> G1	50	7 (13.46)	0.830± 0.029	0.00470	-0.27403	-0.193	
<i>E. granulosus</i> sensu stricto complex	<i>E. granulosus</i> G3	2	2 (3.84)	1± 0.500	0.01351		
Total	52	9			*ND	ND	

*ND: Not determined. * P<0.01. ** P<0.02.

N: Number of isolates; **Nh:** Number of haplotypes; **Hd:** Haplotypes (gene) diversity; **Nd:** Nucleotide diversity.

Table 3: Haplotype distribution of *E. granulosus* G1 and G3 genotypes in human FFPE tissue samples based on partial cytochrome c oxidase subunit 1(cox1) mitochondrial sequence in Mazandaran Province, Northern Iran

Haplotype	Genotype	Accession number
MAZ04	G3	MH920521
MAZ09	G3	MH920524
MAZ02	G1	MH920515
MAZ05	G1	MH920518
MAZ07	G1	MH920520
MAZ11	G1	MH920522
MAZ12	G1	MH920525
MAZ13	G1	MH920526
MAZ15	G1	MH920528

No Indel mutations (deletions/insertion) identified in *E. granulosus* G1 and G3 genotypes. Within the 444bp consensus position of G1 and G3 genotypes, 7 and 5 variable (polymorphic) sites were detected, respectively. The pairwise distance matrix between aligned sequences of G3 genotype

provided an intra-diversity of 1.4% and percent identity of 98.6% while an intra-divergence of 0.3%-1.6% and high percent identity of 98.6%-99.7% were calculated for G1 genotype (Fig. 2). The inter-divergence level between G1 and G3 genotypes was estimated 1.1% to 2.2% (Fig. 2).




Fig. 2: The sequence pairwise distances including intra and inter-divergence levels and percent identity of *E. granulosus* G1 and G3 genotypes among the analyzed sequences

Diversity indices, neutrality indices, and multiple sequence alignment

The diversity indices, neutrality indices, and Hn of G1 and G3 genotypes are given in Table 2. DNA sequence analyses of *E. granulosus* G1 and G3 genotypes indicated high genetic diversity of *Cox1* (G1: Hd: 0.830 and G3: Hd: 1.00). The negative indices of *E. granulosus* G1 isolates (Tajima's D; -0.27403 and Fu's Fs; -0.193), indicating a substantial divergence from neutrality (Table 2). To illustrate the expansion and the divergence time for

unequal population sizes of *Echinococcus*, the indices of Tau: 1.911, Raggedness r (0.0948), and R2 statistic (0.1158) are shown in Fig. 3. The multiple sequence alignments of the nucleotides and amino acids of G1 and G3 genotypes are shown in Fig. 4 and 5. The non-synonymous substitutions of *E. granulosus* G1 occurred at codons 2,3, 85 (MAZ02), 71 (MAZ11), 74 (MAZ13), and 85 (MAZ12), besides two amino acid substitutions were identified at codons 65 and 66 in *E. granulosus* G3 (Fig. 5).




Fig. 3: Expected and observed mismatch distribution for *E. granulosus* G1 genotype inferred by *cox1* gene




Fig. 4: Nucleotide sequence alignment of *cox1* gene based on identified haplotypes of *E. granulosus* G1 and G3 genotypes




Fig. 7: Haplotype network in *E. granulosus* sensu stricto based on mitochondrial sequences of *cox1* from Northern Iran (constructed by PopART software)

Discussion

Findings of this study showed that the human CE with *E. granulosus* G1 and G3 genotypes containing various genetic traits are explicitly being circulated in bordering the Caspian Sea, Northern Iran. Presently, CE/alveolar echinococcosis as an ignored public health concern threat a wide range of at-risk populations, particularly those who settle in hyperendemic areas. Despite the initiate of preventive strategies by Iran's health policymakers, there is an increasing incidence of human hydatidosis in some areas of Iran (e.g., Tabriz city, 10 cystectomy CE cases/1 month) (24–26). Currently, the G1, G2, G3, and G6 strain have been reported from human hydatidosis isolates in Iran (13). Identification and genotyping of *Echinococcus* spp., in human CE isolates are potentially critical to determine probable drug-resistance codons, genetic diversity traits, and transmission patterns of this parasite (27).

In concurrence with earlier studies in Iran, the G1 genotype was the most dominant strain associated

with human hydatidosis. We also detected a considerable haplotype variability of *E. granulosus* G1 (Hd; 0.830) and G3 (Hd; 1.00) genotypes in human clinical isolates obtained from bordering the Caspian Sea, Northern Iran. The occurrence of significant diversity (novel haplotypes) of G1/G3 genotypes indicates that the sheep and buffalo hosts may potentially contribute in maintenance and transmission of probable emergent *Echinococcus* sp., to candies, and will also lead to establishment of resistant alleles to treatment in the region. Furthermore, genetic diversity in *Echinococcus* parasite may frequently take place after high gene flow, in which migration increases the effectual population size in various geographical regions, where the diversity characters are likely dominant (28). In addition, occurring codons 65 and 66 of G3 genotype should be taken into consideration in intensification of *E. granulosus* G3 pathogenicity or development of plausible drug-resistance alleles (21,29). According to the above-mentioned descriptions, our earlier study showed that the gene migration (gene flow) of *cox1* in support of *E. granulosus*. G1 and G3 strains are unambiguously

shared between Northern-Central, Northern-Southeastern, and Northern-Western pair populations in Iran (30-32). Previous study *E. granulosus* isolates from definitive hosts (Dogs and Golden jackals) in Mazandaran northern Iran, using partial *cox1* sequence, 66.7% and 33.3% of isolates belonged to G1 and G3 genotypes, respectively (33). Another study of *E. granulosus* isolates from live-stock and humans in Golestan Province, northern Iran indicated G1 (78.3%), G2 (2.7%), G3 (15%) and G6 (4%) genotypes among the 74 CE isolates (34).

The presence of negative neutrality indices (-0.27403 to -0.193) in support of *E. granulosus* G1 genotype implies to population expansion after the bottleneck incident and purifying selection (32,33). The findings of Tau value (1.911) and unimodal mismatch distribution test revealed that the *E. granulosus* G¹ genotype have newly experienced a population expansion due to mitochondrial *cox1* gene migration of parasite in bordering the Caspian Sea, Northern Iran.

In present study, the majority of *E. granulosus* G3 genotype was only limited to human liver hydatidosis, on the one hand, a letter from Iran shown that, *E. canadensis* G6 isolates (n=8) has a positive tropism to shape the human cerebral CE (34). Although, in the current study, no cases of Central Nervous System CE were obtained, however more samples are needed to confirm the potential tropism of genotypes of *E. granulosus* sensu lato to organ specificity of humans.

Conclusion

In this study, for the first time, G3 genotype (buffalostrain) was identified from two human hydatidosis isolates in Mazandaran Province which indicating the water buffalo can play in the transmission of *Echinococcus* G3 to canids. Present results strengthen our knowledge concerning taxonomic status, transmission patterns of *Echinococcus* parasite to human and heterogeneity aspects of this parasite in Northern Iran which will also assist in

the expansion of strategies for controlling of intermediate hosts and monitor of clinical isolates in the region.

Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Acknowledgements

This research is a part of the first author's Ph.D. thesis. The authors thank all colleagues working in Toxoplasmosis Research Centre (TRC) at Mazandaran University of Medical Sciences. This work was supported by a grant (No. 2929) from the Deputy of Research, Mazandaran University of Medical Sciences, Sari, Iran.

Conflict of interest

The authors declare that they have no conflict of interest.

References

1. Agudelo Higueta NI, Brunetti E, McCloskey C (2016). Cystic Echinococcosis. *J Clin Microbiol*, 54 (3):518-523.
2. Otero-Abad B, Torgerson PR (2013). A systematic review of the epidemiology of echinococcosis in domestic and wild animals. *PLoS Negl Trop Dis*, 7 (6):e2249.
3. Brunetti E, Garcia HH, Junghans T, et al (2011). Cystic echinococcosis: chronic, complex, and still neglected. *PLoS Negl Trop Dis*, 5 (7):e1146.
4. Zhang W, Zhang Z, Wu W, et al (2015). Epidemiology and control of echinococcosis in central Asia, with particular reference to the People's Republic of China. *Acta Trop*, 141 (Pt B):235-43.
5. Carmena D, Sanchez-Serrano LP, Barbero-Martinez I (2008). *Echinococcus granulosus* infection in Spain. *Zoonoses Public Health*, 55 (3):156-65.

6. Craig PS, McManus DP, Lightowlers MW, et al (2007). Prevention and control of cystic echinococcosis. *Lancet Infect Dis*, 7 (6):385-94.
7. Buishi I, Njoroge E, Zeyhle E, et al (2006). Canine echinococcosis in Turkana (north-western Kenya): a coproantigen survey in the previous hydatid-control area and an analysis of risk factors. *Ann Trop Med Parasitol*, 100 (7):601-10.
8. Siracusano A, Delunardo F, Teggi A, Ortona E (2012). Host-parasite relationship in cystic echinococcosis: an evolving story. *Clin Dev Immunol*, 2012:639362.
9. Buishi IE, Njoroge EM, Bouamra O, Craig PS (2005). Canine echinococcosis in northwest Libya: assessment of coproantigen ELISA, and a survey of infection with analysis of risk-factors. *Vet Parasitol*, 130 (3-4):223-232.
10. Buishi I, Walters T, Guildea Z, et al (2005). Reemergence of canine *Echinococcus granulosus* infection, Wales. *Emerg Infect Dis*, 11 (4):568-571.
11. Belli S, Akbulut S, Erbay G, Kocer NE (2014). Spontaneous giant splenic hydatid cyst rupture causing fatal anaphylactic shock: a case report and brief literature review. *Turk J Gastroenterol*, 25 (1):88-91.
12. Engin G, Acunas B, Rozanes I, Acunas G (2000). Hydatid disease with unusual localization. *Eur Radiol*, 10 (12):1904-1912.
13. Galeh TM, Spotin A, Mahami-Oskouei M, et al (2018). The seroprevalence rate and population genetic structure of human cystic echinococcosis in the Middle East: A systematic review and meta-analysis. *Int J Surg*, 51:39-48.
14. Ebrahimipour M, Sadjjadi SM, Yousofi Darani H, Najjari M (2017). Molecular Studies on Cystic Echinococcosis of Camel (*Camelus dromedarius*) and Report of *Echinococcus ortleppi* in Iran. *Iran J Parasitol*, 12 (3):323-331
15. Fadakar B, Tabatabaei N, Borji H, Naghibi A (2015). Genotyping of *Echinococcus granulosus* from goats and sheep indicating G7 genotype in goats in the Northeast of Iran. *Vet Parasitol*, 214 (1-2):204-7.
16. Fasihi Harandi M, Budke CM, Rostami S (2012). The monetary burden of cystic echinococcosis in Iran. *PLoS Negl Trop Dis*, 6 (11):e1915.
17. McManus DP, Thompson RC (2003). Molecular epidemiology of cystic echinococcosis. *Parasitology*, 127:S37-51
18. McManus DP (2002). The molecular epidemiology of *Echinococcus granulosus* and cystic hydatid disease. *Trans R Soc Trop Med Hyg*, 96 Suppl 1:S151-7.
19. Gholami I, Daryani A, Sharif M, et al (2011). Seroepidemiological survey of helminthic parasites of stray dogs in Sari City, northern Iran. *Pak J Biol Sci*, 14 (2):133-7
20. Schneider R, Gollackner B, Schindl M, et al (2010). *Echinococcus canadensis* G7 (pig strain): an underestimated cause of cystic echinococcosis in Austria. *Am J Trop Med Hyg*, 82 (5):871-874.
21. Bowles J, Blair D, McManus DP (1992). Genetic variants within the genus *Echinococcus* identified by mitochondrial DNA sequencing. *Mol Biochem Parasitol*, 54(2):165-73.
22. Bandelt HJ, Forster P, Rohl A (1999). Median-joining networks for inferring intraspecific phylogenies. *Mol Biol Evol*, 16 (1):37-48.
23. Rozas J, Sanchez-DelBarrio JC, Messeguer X, Rozas R (2003). DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics*, 19 (18):2496-7.
24. Noori J, Spotin A, Ahmadpour E, et al (2018). The potential role of toll-like receptor 4 Asp299Gly polymorphism and its association with recurrent cystic echinococcosis in postoperative patients. *Parasitol Res*, 117 (6):1717-1727.
25. Grosso G, Gruttadauria S, Biondi A, et al (2012). Worldwide epidemiology of liver hydatidosis including the Mediterranean area. *World J Gastroenterol*, 18 (13):1425-1437.
26. Woolhouse ME, Gowtage-Sequeria S (2005). Host range and emerging and reemerging pathogens. *Emerg Infect Dis*, 11 (12):1842-1847.
27. Rojas CAA, Romig T, Lightowlers MW (2014). *Echinococcus granulosus* sensu lato genotypes infecting humans—review of current knowledge. *Int J Parasitol*, 44 (1):9-18
28. Mahami-Oskouei M, Kaseb-Yazdanparast A, Spotin A, et al (2016). Gene flow for *Echinococcus granulosus* metapopulations determined by mitochondrial sequences: a reliable approach for reflecting epidemiological drift of parasite among neighboring countries. *Exp Parasitol*, 171:77-83
29. Mokhtari Amirmajdi M, Sankian M, Eftekharzadeh Mashhadi I, et al(2011). Apoptosis of human lymphocytes after

- exposure to hydatid fluid. *Iran J Parasitol*, 6 (2):9-16
30. Spotin A, Mahami-Oskouei M, Harandi MF, et al (2017). Genetic variability of *Echinococcus granulosus* complex in various geographical populations of Iran inferred by mitochondrial DNA sequences. *Acta Trop*, 165:10-16
 31. Gholami S, Sosari M, Fakhar M, et al (2012). Molecular Characterization of *Echinococcus granulosus* from Hydatid Cysts Isolated from Human and Animals in Golestan Province, North of Iran. *Iran J Parasitol*, 7 (4):8-16
 32. Sadjjadi S, Mikaeili F, Karamian M, et al (2013). Evidence that the *Echinococcus granulosus* G6 genotype has an affinity for the brain in humans. *Int J Parasitol*, 43 (11):875-7.
 33. Gholami S, Jahandar H, Abastabar M, et al (2016). *Echinococcus granulosus* sensu stricto in dogs and jackals from Caspian Sea region, northern Iran. *Iran J Parasitol*, 11(2):186-194.
 34. Sharbatkhori M, Tanzifi A, Rostami S, et al (2016). *Echinococcus granulosus* sensu lato genotypes in domestic livestock and humans in Golestan province, Iran. *Rev Inst Med Trop Sao Paulo*, 58:38.