



## Seroepidemiology of Herpes Simplex Virus Type 1 and 2 in Northern Iran

*S Rezaei-Chaparpordi<sup>1</sup>, \*M Assmar<sup>1</sup>, N Amirmozafari<sup>2</sup>, L Modiri<sup>1</sup>, A Massiha<sup>1</sup>, S Shokri-Fashtali<sup>3</sup>, Z Gholizadeh<sup>4</sup>, S Akbari<sup>1</sup>*

1. *Dept. of Microbiology, Labijan Islamic Azad University, Labijan, Iran*
2. *Dept. of Microbiology, Tehran University of Medical Sciences, Tehran, Iran*
3. *Dept. of Mathematics, Labijan Islamic Azad University, Labijan, Iran*
4. *Zist Faravard Pars Company, Rasht, Iran*

(Received 28 Oct 2011; accepted 12 Jul 2012)

### Abstract

**Background:** Herpes simplex virus (HSV) type 1 and 2 are common infectious agents worldwide. Data on prevalence of HSV-1 and HSV-2 infection are limited in Asia, especially in Iran. The aim of this study was to determine the seroprevalence of HSV type 1 and 2 based on age, gender, marital status, education, living area, job, symptoms and history of disease variables.

**Methods:** The study population included 800 randomly selected persons from laboratories in Gilan Province, Iran, from 2010 to 2011. Demographic data gathered by a well-designed questionnaire and for serological studies, blood samples were collected and centrifuged. ELISA HSV-1, 2 and HSV-2 specific ELISA kits were used to determine IgG type specific antibodies in sera samples. Person's chi-square test was applied to compare HSV-1 and HSV-2 seropositivities.

**Results:** HSV-1 and HSV-2 IgG antibodies were positive in 467 (58.4%) and 28 (3.5%) subjects, respectively. There was significant correlation between age, marital status, job, symptoms, history of disease and HSV seroprevalence ( $P < 0.05$ ).

**Conclusion:** Our findings were in agreement with prior studies in which HSV-1 infections was more prevalent than HSV-2 and seropositivity increased with age.

**Keywords:** HSV-1, HSV-2, ELISA, Prevalence, Iran

### Introduction

Herpes simplex virus (HSV) infections are caused by two types of viruses, type 1 (HSV-1) and type 2 (HSV-2), that are among the most common human viral infections worldwide (1-3). HSV-1 is transmitted by two ways: 1-casual contacts during childhood, 2-sexual contacts, but HSV-2 is a virus that is mostly transmitted sexually (4). HSV can cause severe systematic neonatal infection by transmission before or during birth (5, 6). HSV-2 is of public health importance as one of the commonest causes of genital ulceration worldwide and

implicated as an important co-factor for HIV infection" (7). HSV-1 was associated predominantly with orolabial ulceration; however, recent changes in HSV-1 and HSV-2 epidemiology have been reported, with an increase in genital and neonatal herpes particularly caused by HSV-1 (8-14). However, routine surveillance is hindered as many of those infected remain asymptomatic or fail to present to health services (15). The recent commercial development of type specific enzyme immunoassays (ELISA) that reliably distinguish be-

tween antibodies to HSV-1 and HSV-2, enable serological studies which can measure both symptomatic and asymptomatic infection. Seropositivity is a potential indicator of infectivity and can be used to guide and reform behavioral patterns for prevention of HSV transmission (16). With several HSV candidate vaccines in clinical trial and antiviral therapy available, a clear understanding of epidemiology of HSV in different populations is required to develop the most effective and appropriate preventive and control strategies (7). The present study was the first study on determination of seroepidemiology of HSV infection in Gilan Province, northern Iran.

### Materials & Methods

Using cluster sampling method, 800 patients were selected randomly from Rasht, Anzali, Lahijan, Somea Sara and Hashtpar laboratories in Gilan Province, Iran, from 2010 to 2011.

After obtaining a verbal consent, their demographic data were recorded via a well designed questionnaire. Questionnaire included questions about age, sex, marital status, level of education, living area, job, clinical symptoms and history of disease. Having explained our goal, 5ml of peripheral venous blood was obtained, then allowed to clot and centrifuged at 1000×g for 10 minutes. Sera were collected into microtubes and transferred to the laboratory in cool box. Sera samples were frozen and stored at -20°C. Type specific serum antibodies to HSV-1 and HSV-2 were detected using HSV-1,2 and HSV-2 ELISA IgG kits (Vircell Spanish kits). ELISA tests were performed according to the manufacturer's instructions.

Person's chi-square test was applied to compare HSV-1 and HSV-2 seropositivities among different groups of randomly selected persons. Statistical significance was set at  $P < 0.05$ .

### Results

The study group comprised subjects aged from 1-85 years old. HSV infections were positive in 61.9%

(495) of persons. HSV-1 and HSV-2 seropositivity was 58.4% (467) and 3.5% (28), respectively.

As expected, HSV-1 seroprevalence was significantly higher than that of HSV-2. Patients were classified to three age groups: 0-19, 20-39, >40 years old. The most HSV prevalence belonged to >40 (74.6%) years old persons.

The study population included 306 (60.8%) males and 494 (62.6%) females (Table 1). Seropositivity in persons who were married (71.3%) and lived in villages (70.6%) was more than single (33.7%) persons that lived in city (60.5%).

**Table 1:** Prevalence of HSV antibodies in the sample studied by age, sex, marital status, living area and education level

Variables	Positive n (%)	Negative n (%)
<b>Age</b> ( $P= 0$ )		
0-19	42(28.6)	105(71.4)
20-39	180(62.7)	107(37.3)
>40	273(74.6)	93(25.4)
<b>Sex</b> ( $P=0.653$ )	186(60.8)	120(39.2)
Males		
Females	309(62.6)	185(37.4)
<b>Marital status</b> ( $P= 0$ )		
Single	67(33.7)	132(67.3)
Married	428(71.2)	173(28.8)
<b>Living area</b> ( $P=0.064$ )		
City	418(60.5)	273(39.5)
Village	77(70.6)	32(29.4)
<b>Education level</b> ( $P=0.271$ )		
University degree	83(53)	60(42)
High school degree	164(62.4)	99(37.6)
Guidance school degree	70(61.4)	44(38.6)
Elementary school degree	80(70.2)	34(29.8)
Illiterate	98(59)	68(41)

According to education level, we classified patients to 5 groups. Results showed high prevalence rate in elementary educated persons (Table1). There were patients with different kinds of jobs but seropositivity was high among the barbers and beauticians (100%).

Some of the subjects had symptoms that were classified into 4 groups as follows:1- general

symptoms (69.8%) including fatigue, backache, headache, influenza like symptoms 2- sore and eczema, 3-pain during urination, 4- Alzheimer's disease. Seroprevalence was 87.8% in persons with Alzheimer's disease. Persons who had history of genital, skin and oral diseases were more seropositive (73.9%) than others (Table 2). According to chi-square test there was significant correlation between age, marital status, job, symptoms and history of disease with seropositivity ( $P < 0.05$ ).

**Table 2:** Prevalence of HSV antibodies in the sample studied by job, clinical symptoms and history of disease

Variables	Positive n(%)	Negative n(%)
<b>Job</b> ( $P= 0$ )	5(62.5)	3(37.5)
Physician and other related job		
Barber	2(100)	0(0)
Farmer	16(88.9)	1(11.1)
Employee	42(70)	18(30)
Teacher	15(75)	5(25)
Worker	9(90)	1(10)
Free job	64(65.3)	34(34.7)
Other	341(58.6)	241(41.4)
<b>Clinical symptoms</b> ( $P= 0.001$ )		
General symptoms	199(69.8)	86(30.2)
Eczema & sore	34(82.9)	7(17.1)
Pain during urinate	43(76.8)	13(23.2)
Alzheimer disease	36(87.8)	5(12.2)
Without clinical symptoms	258(55.5)	207(44.5)
<b>History of disease</b> ( $P=0$ )		
With history	289(73.9)	102(26.1)
Without history	206(50.4)	203(49.6)

## Discussion

Although the incidence and prevalence of herpes simplex infection is growing, there is little information about HSV prevalence for Iranian population that shows the necessity of more studies on this subject to be done.

This paper is the first to our knowledge to present the comparative seroepidemiology of HSV-1 and HSV-2 in Gilan Province. The seroprevalence of HSV-1 and HSV-2 IgG antibody was 58.4% and 3.5%, respectively. Like other studies, we found

that HSV-1 is more prevalent than HSV-2 and seroprevalence of HSV infection increased with age. The most prevalence rate was found in old and married persons (>40) probably because they have more time for contact to receive the virus.

Barbers are always in contact with skin and customer's saliva so we expected to have high prevalence in this group of people.

The seroprevalence was high in persons with Alzheimer's disease and history of oral, skin, genital diseases.

"Up to 70% of genital HSV infections are unrecognized, hence seroepidemiological studies are critical for understanding the pattern and distribution of infection within populations" (17). According to Looker and Xu studies in United states, Europe and some developed countries, HSV-1 is the main causative agent of genital herpes (18,19). In a study in Europe, HSV-1 was 52% (Finland), 57% (Netherlands), 67% (Belgium), 81% (Czech republic), 84% (Bulgaria) and HSV-2 seroprevalence was 24% (Bulgaria), 14% (Germany), 13% (Finland), 11% (Belgium), 9% (Netherland), 6% (Czech republic), and 4% (England and Wales) (7). In Brazil, HSV-1 and HSV-2 seroprevalence was 67.2% and 11.3%, respectively. In this study, seroprevalence increased with age and past history of STD was a major risk factor for HSV-2 infection (20).

There is a correlation between educational level and HSV seroprevalence. HSV-1 and HSV-2 seroprevalence rate was 80% and 19.3% (21).

Another study in USA showed declines in HSV-1 (57.7%) and HSV-2 (17%) seroprevalence rate during 1999-2004 (19). The seroprevalence of HSV-1 and HSV-2 in our study was lower than some African countries where its HSV-1 and HSV-2 seroprevalence rate has been reported 90% and 82%, respectively (22).

This study highlights the frequency of HSV infection in Gilan province, Iran. Reducing risk factors and awareness of the population are among the helpful health policies against this infection.

## Ethical considerations

Ethical issues (Including plagiarism, Informed Consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc) have been completely observed by the authors.

## Acknowledgment

Sera samples for this study were gathered from laboratories in Gilan Province. This study has been done by personal expenditure. Authors wish to thank Dr. Afrah and Dr. Mehrvarz in Afrah laboratory, Dr. Habibzadeh, Dr. Shahbazi and Mr. Ahmadi in Razi laboratory, Dr. Omidvari in Omidvari laboratory, Dr. Movahed in Mehr laboratory, Dr. Sharifi in Sharif laboratory, Dr. Aghae in Salamat laboratory, Noorani hospital staff and Mrs. Shahidi in Pars laboratory staff. The authors declare that there is no conflict of interest.

## References

- Nahmias AJ, Lee FK, Beckmann-Nahmias F (1990). Seroepidemiological and sociological patterns of herpes simplex infections in the world. *Scand J Infect Dis Suppl*,69:19-36.
- Schomogyi M, Wald A, Corey L (1998). Herpes simplex virus-2 infections: an emerging disease? *Infect Dis Clin North Am*, 12(1):47-61.
- Kasubi MJ, Nilsen A, Marsden HS, Bergström T, Langeland N, Haarr L, et al. (2006). Prevalence of antibodies against herpes simplex virus types 1 and 2 in children and young people in an urban region in Tanzania. *J Clin Microbiol*, 44(8):2801-7.
- Whitley RC (2001). Herpes simplex viruses. In: *In Fields virology*. Ed, Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA, Roizman B, Straus SE. Philadelphia, New York, Lippincott Williams & Wilkins, pp 2461- 2509.
- Tookey P, Peckham CS (1996). Neonatal herpes simplex virus infection in the British Isles. *Paediatr Perinat Epidemiol*,10:432-42.
- Brown ZA, Selke S, Zeh J, et al. (1997). The acquisition of herpes simplex virus during pregnancy. *N Engl J Med*, 337:509-15.
- Pebody RG, Andrews N, Brown D, Gopal R, De Melker H, François G, et al. (2004). The seroepidemiology of herpes simplex virus type 1 and 2 in Europe. *Sex Transm Infect*,80(3):185-91.
- Lafferty WE, Downey L, Celum C, et al. (2000). Herpes simplex type 1 as a cause of genital herpes: impact on surveillance and prevention. *J Infect Dis*,181:1454-7.
- Langenberg AG, Corey L, Ashley RL, et al. (1999). A prospective study of new infections with herpes simplex virus type 1 and type 2: Chiron HSV Vaccine Study Group. *N Engl J Med*, 341:1432-8.
- Ades A, Peckham C, Dale G, et al. (1989). Prevalence of antibodies to herpes simplex virus types 1 and 2 in pregnant Women, and estimated rates of infections. *J Epidemiol Commun Health*, 43: 53-60.
- Ross J, Smith I, Elton R (1993). The epidemiology of herpes simplex types 1 and 2 infection of the genital tract in Edinburgh 1978-91. *Genitourin Med*, 69:381-3.
- Cowan F, Johnson A, Ashley R (1996). Relationship between antibodies to herpes simplex virus (HSV) and symptoms of HSV infection. *J Infect Dis*, 174:470-5.
- Lowhagen GB, Tunback P, Andersson K, et al. (2000). First episodes of genital herpes in a Swedish STD population: a study of epidemiology and transmission by the use of herpes simplex (HSV) typing and specific serology. *Sex Transm Infect*,76:179-82.
- Gaytant M, Steegers EA, van Cromvoirt P, et al(2000). Incidence of herpes neonatorum in Netherlands. *Ned Tijdschr Geneeskde*,144:1832-6.
- Nahmias AJ, Lee FK, Beckman-Nahmias S, et al. (1990). Seroepidemiological and sociological patterns of herpes simplex infection in the world. *Scand J Infect Dis Suppl*, 69:19-36.
- Levett PN (2005). Seroprevalence of HSV-1 and HSV-2 in Barbados. *Med Microbiol Immunol*,194(1-2): 105-7.
- Cowan FM, French RS, Mayaud P, Gopal R, Robinson N, de Oliveira SA, et al. (2003). Seroepidemiological of herpes simplex virus type 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka. *Sex Transm Infect*, 79(4): 286-90.

18. Looker KJ, Garnett GP (2005). A systematic review of the epidemiology and interaction of herpes simplex virus types 1 and 2. *Sex Transm Infect*, 81(2):103-7.
19. Xu F, Sternberg MR, Kottiri BJ, McQuillan GM, Lee FK, Nahmias AJ, et al. (2006). Trends in herpes simplex virus type 1 and type 2 seroprevalence in the United States. *JAMA*, 296 (8): 964-73.
20. Clemens SC, Farhat CK (2010). Seroprevalence of herpes simplex 1-2 antibodies in Brazil. *Rev Saude Publica*, 44 (4): 1-8.
21. Bünzli D, Wietlisbach V, Barazzoni F, Sahli R, Meylan PR (2004). Seroepidemiology of Herpes Simplex virus type 1 and 2 in Western and Southern Switzerland in adults aged 25-74 in 1992-93: a populationbased study. *BMC Infect Dis*, 17:4-10.
22. Smith JS, Robinson NJ (2002). Age-specific prevalence of infection with herpes simplex virus types 2 and 1: a global review. *J Infect Dis*, 186:3-28.