



## **Comparison of Standardized Plant Analysis Risk Human Reliability Analysis (SPAR-H) and Cognitive Reliability Error Analysis Methods (CREAM) in Quantifying Human Error in Nursing Practice**

***Iraj MOHAMMADFAM<sup>1</sup>, Maryam MOVAFAGH<sup>1</sup>, \*Saeid BASHIRIAN<sup>2</sup>***

1. *Dept. of Occupational Hygiene, School of Public Health and Research Center for Health Sciences, Hamadan University of Medical Sciences, Hamadan, Iran*
2. *Dept. of Health Education, Hamadan University of Medical Sciences, Hamadan, Iran*

**\*Corresponding Author:** Email: bashirian@umsha.ac.ir

**(Received 03 Oct 2015; accepted 04 Nov 2015)**

### **Dear Editor-in-Chief**

Nurses spend much more time dealing with patients compared with other healthcare personnel. This group has a significant role in identification of threatening complexities of patients, removing medical errors and in keeping patients safe, on the other hand, a considerable amount of human errors is committed by nurses (1). Therefore, nurses have to face with many occupational risks compared with other jobs and may suits relating to occupational negligence (1). Human errors associated with nursing duties; in particular, medication error has been identified as a central concern for the nursing profession (1). As the result of such events, patient is hurt unwantedly and staff suffers a great deal of damage. These events often result from the change of systemic specifications in duties, group, work environment, and organization (3).

The present study was a cross sectional research aimed at comparing two human error identification and assessment methods, SPAR-H and CREAM, in nursing practice at a hospital in Iran. We investigated 31 duties according to job description announced by Iran Nursing Organization and professors of nursing.

In the first step, we analyzed occupational duties using Hierarchical Task Analysis (HTA) method (4). Next, different tasks are executed for each sub task.

After completing CREAM and SPAR-H worksheets, we first grouped the worksheets for each duty and analyzed total error probability and error risk using SPSS-16 (Chicago, IL, USA). In order to compare the two techniques based on multivariable decision making, we used analytical hierarchy process (5).

The findings of demonstrated that, the estimated human error probability by SPAR-H was higher than those estimated by CREAM, but the most critical task, "taking initial measures for cardiopulmonary resuscitation until the arrival of resuscitation team", was the same in both techniques. Moreover, it was determined that both techniques have similar results in identifying and assessing critical tasks with high probability of human error (Table 1). Furthermore, there was more difference between these two techniques in ranking potential errors with low probabilities; for instance, preparing bed before arrival of patient have the rank of 27 when we used SPAR-H technique 27, while its rank using CREAM technique was determined at 10 (Table 1).

Based on previous studies in this field and also eliciting domain expert knowledge, seven criteria were selected for comparing two techniques with each other (6).

**Table 1:** Human Error Probability (HEP) and CFPt for each duty nursing using SPAR-H and CREAM methods

| Duties  | HEP (SPAR-H) | Rank (SPAR-H) | CFPt* (CREAM) | Rank (CREAM) |
|---|--------------|---------------|---------------|--------------|
| Taking initial measures for cardiopulmonary resuscitation until the arrival of resuscitation team | 0.37180      | 1             | 0.0591        | 1            |
| Notification of code and starting initial steps of cardiopulmonary resuscitation                  | 0.24489      | 2             | 0.02086       | 4            |
| Adjusting and using DC shock in emergency cases   | 0.23295      | 3             | 0.02108       | 2            |
| Performing intubation in emergency cases  | 0.19574      | 4             | 0.01730       | 6            |
| Taking urgent actions for dysrhythmia   | 0.12217      | 5             | 0.01189       | 11           |
| Recording data in integrated hospital system  | 0.08552      | 6             | 0.02088       | 3            |
| Heart monitoring  | 0.05340      | 7             | 0.01413       | 9            |
| Performing suction of airway secretions and tracheal tube   | 0.04886      | 8             | 0.00888       | 15           |
| Education before, during and after hospitalization  | 0.04596      | 9             | 0.02060       | 5            |
| Establishing IV line  | 0.04415      | 10            | 0.00489       | 24           |

\*Cognitive Failure Probability total

These criteria are as follows:

1. Training courses required for implementation (TN)
2. Cost of Execution (CE)
3. Duration of Execution (DE)
4. Focus on Social Factors(SF)
5. Focus on Organizational Factors(OF)
6. Focus on Individual Factors(IF)
7. Preventive Measures (PM).

Table 2 shows results of the final weight of each component and criteria based on AHP method for both SPAR-H and CREAM methods.

The final weight in SPAR-H and CREAM methods is calculated as follows:

$$\text{SPAR-H} = 0.967 \times 0.834 + 2.834 \times 0.166 + 6.2 \times 0.33 + 4.12 \times 0.834 + 1.192 \times 0.166 + 2.567 \times 0.88 + 0.387 \times 0.75 = 8.094234$$

$$\text{CREAM} = 0.967 \times 0.166 + 2.834 \times 0.834 + 6.2 \times 0.66 + 4.12 \times 0.166 + 1.192 \times 0.834 + 2.567 \times 0.66 + 0.387 \times 0.25 = 10.0851$$

Final score in CREAM (10.0851) was more than SPAR-H (8.094234) method, then, it can be concluded that CREAM is more efficient than SPAR-H.

**Table 2:** Final weight of the components and criteria's in SPAR-H and CREAM methods

|              | DE    | TN    | IF   | OF    | SF    | PM    | CE    |
|--------------|-------|-------|------|-------|-------|-------|-------|
| SPAR-H       | 0.834 | 0.166 | 0.33 | 0.834 | 0.166 | 0.33  | 0.75  |
| CREAM        | 0.166 | 0.834 | 0.66 | 0.166 | 0.834 | 0.66  | 0.25  |
| Final weight | 0.967 | 2.834 | 6.2  | 4.12  | 1.192 | 2.567 | 0.387 |

## Acknowledgments

This research has the financial support of Deputy of Research, Hamadan University of Medical Sciences, Iran. The authors declare that there is no conflict of interest.

## References

1. Parry AM, Barriball KI, While AE (2015). Factors contributing to Registered Nurse medication administration error: A narrative review. *Int J Nurs Stud*, 52(1), 403-420.
2. Donchin Y, Gopher D, Olin M, et al. (1995). A look into the nature and causes of human errors in the intensive care unit. *Crit Care Med*, 23(2):294-300.
3. McNutt RA, Abrams R, Arons DC (2002). Patient safety efforts should focus on medical errors. *JAMA*, 287(15): 1997-2001.
4. Stanton NA (2006). Hierarchical task analysis: Developments, applications, and extensions. *Appl Ergon*, 37(1): 55-79.
5. Caputo AC, Pelagagge PM, Salini P (2013). AHP-based methodology for selecting safety devices of industrial machinery. *Safety Sci*, 53: 202-18.
6. Rouvroye JL, van den Bliek EG (2002). Comparing safety analysis techniques. *Rehabil Eng Syst Safe*, 75(3): 289-94.