

Letter to the Editor

Clinical Problem: Cancer Biopsy

*Mohammad Reza SASANI

Dept. of Radiology, Medical Imaging Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

*Correspondence: Email: sasanimr@sums.ac.ir

(Received 09 Feb 2016; accepted 24 Mar 2016)

Dear Editor-in-Chief

A letter to the editor entitled "Cancer Biopsy in Ancient Persian Medicine: Do or Not to Do!" was published in Iran J Public Health (1). I would like to focus attention on authors' suggestion for using non-invasive methods such as laboratory parameters and biomarkers, for tumor detection. Advancements in oncology resulted in development of different protocols for cancer treatment. Accordingly, accurate diagnosis base on pathologic specimen is very important particularly to select a suitable treatment protocol and therefore, doing tumor biopsy is inevitable. Although concern for tumor seeding during biopsy still exists but several studies, for instance (2-4) have found that, its likelihood is very low. Furthermore, advantages of correct cancer diagnosis is would usually weigh more than potential risks of biopsy (3, 4).

References

- Gorji N, Moeini R, Pasalar P et al. (2015). Cancer Biopsy in Ancient Persian Medicine: Do or Not to Do! *Iran J Public Health*, 44(10): 1440– 1441.
- Loughran CF, Keeling CR (2011). Seeding of tumour cells following breast biopsy: a literature review. Br J Radiol, 84(1006):869-874.
- 3. Robertson EG, Baxter G (2011). Tumour seeding following percutaneous needle biopsy: the real story! *Clin Radiol*, 66(11):1007-1014.
- Volanis D, Neal DE, Warren AY, Gnanapragasam VJ (2015). Incidence of needle-tract seeding following prostate biopsy for suspected cancer: a review of the literature. *BJU Int*, 115(5):698-704.